

ECE-597: Probability, Random Processes, and Estimation
Homework # 6

Due: Tuesday April 26, 2016

Exam 2, Friday May 6

From the textbook: 8.24, 8.26 (a,b), 8.34, 8.35, 8.37, 8.42

Hints and Answers:

$$8.24, K_{\mathbf{YY}}[n] = \sigma^2 \left\{ (1 + |\alpha|^2) \rho^{|n|} - \alpha^* \rho^{|n+1|} - \alpha \rho^{|n-1|} \right\}, \alpha = \rho \text{ or } \alpha = 1/\rho$$

$$8.26, \mu_{\mathbf{y}}[n] = \frac{5}{1-\rho}, E[y[n]^2] = \frac{25}{(1-\rho)^2} + \frac{\sigma_w^2}{1-\rho^2}$$

$$8.34, h[m] = K_{\mathbf{XW}}[m], S_{\mathbf{XW}}(\omega) = H(\omega)$$

$$8.35, S_{\mathbf{XY}}(\omega) = H^*(\omega)S_{\mathbf{XX}}(\omega)$$

$$8.37, S_{\mathbf{XW}}(\omega) = \frac{S_{\mathbf{XX}}(\omega)}{\sqrt{S_{\mathbf{YY}}(\omega)}}, h[n] = R_{\mathbf{XW}}[n] \text{ for } n = 0, 1, \dots, N-1$$

$$8.42, \mu_{\mathbf{X}}[n] = \frac{n}{2}(s_1 - s_2), R_{\mathbf{XX}}[n_1, n_2] = \frac{\min(n_1, n_2)}{4}(s_1 + s_2)^2 + \frac{n_1 n_2}{4}(s_1 - s_2)^2$$

Additional Problem

a) We can write $a^{|m|}$ as

$$a^{|m|} = a^m u[m-1] + a^{-m} u[-m]$$

Using this relationship derive the z - transform pair

$$a^{|m|} \leftrightarrow \frac{1-a^2}{(1-az^{-1})(1-az)}$$

b) Assume we have an autoregressive model

$$x[n] = ax[n-1] + v[n]$$

with $|a| < 1$, $v[n]$ zero mean white noise. Show that

$$\begin{aligned} S_{\mathbf{XX}}(z) &= \frac{\sigma_v^2}{(1-az^{-1})(1-az)} \\ R_{\mathbf{XX}}[m] &= \frac{\sigma_v^2 a^{|m|}}{1-a^2} \end{aligned}$$

c) Assume we have a moving average process

$$x[n] = v[n] - bv[n-1]$$

with $|b| < 1$, and $v[n]$ zero mean white noise. Show that

$$S_{\mathbf{XX}}(z) = \sigma_v^2 (1-bz^{-1})(1-bz)$$

$$R_{\mathbf{XX}}[m] = (1+b^2)\sigma_v^2 \delta[m] - b\sigma_v^2 \delta[m+1] - b\sigma_v^2 \delta[m-1]$$

d) Assume we have a simple autoregressive moving average model

$$x[n] = ax[n - 1] - bv[n - 1] + v[n]$$

with $|a| < 1, |b| < 1$. Show that

$$S_{\mathbf{XX}}(z) = \sigma_v^2 \frac{(1 - bz^{-1})(1 - bz)}{(1 - az^{-1})(1 - az)}$$

and find $R_{\mathbf{XX}}[m]$