

**Composition of Foods  
Raw, Processed, Prepared  
USDA National Nutrient Database for Standard  
Reference, Release 19**

August 2006

U.S. Department of Agriculture  
Agricultural Research Service  
Beltsville Human Nutrition Research Center  
Nutrient Data Laboratory  
10300 Baltimore Avenue  
Building 005, Room 107, BARC-West  
Beltsville, Maryland 20705

U.S. Department of Agriculture, Agricultural Research Service, USDA Nutrient Data Laboratory. 2006. USDA National Nutrient Database for Standard Reference, Release 19.

USDA Nutrient Data Laboratory website: <http://www.ars.usda.gov/nutrientdata>

Mention of trade names, commercial products, or companies in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture over others not mentioned.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Issued August 2006

## Contents

Introduction.....	1
Specific Changes for SR19 .....	1
Data Files .....	2
Reports .....	3
File Content.....	3
Food Description File.....	3
Nutrients.....	4
Proximates.....	8
Minerals .....	10
Vitamins.....	10
Lipid Components.....	14
Amino Acids.....	18
Weights and Measures .....	19
Footnotes.....	19
Sources of Data .....	19
Explanation of File Formats .....	20
Relational Files .....	20
Food Description File .....	21
Food Group Description File.....	23
Nutrient Data File.....	23
Nutrient Definition File.....	24
Source Code File.....	25
Data Derivation Code Description File .....	25
Weight File.....	26
Footnote File.....	27
Sources of Data File.....	27
Sources of Data Link File.....	28
Abbreviated File .....	28
Update Files .....	31
Summary.....	33
References.....	34
Appendix A. Abbreviations Used in Short Descriptions.....	A-1
Appendix B. Other Abbreviations .....	B-1

## **Introduction**

The USDA National Nutrient Database for Standard Reference (SR) is the major source of food composition data in the United States. It provides the foundation for most food composition databases in the public and private sectors. As information is updated, new versions of the database are released. This version, Release 19 (SR19), contains data on 7,293 food items and up to 140 food components. It replaces SR18 issued in August 2005.

SR19 includes composition data for all the food groups and nutrients published in the 21 volumes of "Agriculture Handbook 8" (U.S. Department of Agriculture 1976–92), and its four supplements (U.S. Department of Agriculture 1990–93), which superseded the 1963 edition (Watt and Merrill, 1963). Since 1992, updated data have been published electronically on the USDA Nutrient Data Laboratory (NDL) website. SR19 supersedes all previous releases, including the printed versions, in the event of any differences.

With the first release of data from our new Nutrient Databank System (NDBS) in July 2001, we changed some formats and added fields to improve the descriptive information for food items and the statistical information about the nutrient values. While data in previous releases have been moved to the new NDBS, they may not have been updated through the complete system. Therefore, many of these new fields contain data only for those items that have been processed through the new NDBS and it will take a number of years before they are populated for most food items in the database.

We compiled data from published and unpublished sources. Published sources include the scientific and technical literature. Unpublished data include those obtained from the food industry, other government agencies, and research conducted under contracts initiated by USDA Agricultural Research Service (ARS). These analyses are currently conducted under the National Food and Nutrient Analysis Program (NFNAP), in cooperation with the National Cancer Institute and 16 other offices and institutes of the National Institutes of Health (Pehrsson et al., 2000). Data from the food industry represents the nutrient content of a specific food or food product at the time the data is sent to NDL. The values may change due to reformulations or other processing changes by individual companies between the time that SR is released and the next update of the database. Values in the database may be based on the results of laboratory analyses or calculated by using appropriate algorithms, factors, or recipes, as indicated by the source code in the Nutrient Data file. Not every food item may contain a complete nutrient profile.

### **Specific Changes for SR19**

We have made several major changes to the database since the last release, as listed below:

- Included nutrient values for about 500 food items for fluoride from the USDA National Fluoride Database of Selected Beverages and Foods - Release 2 (USDA, 2005). Some values, such as regional values, were not incorporated into SR.

- Included nutrient values for total choline and betaine for approximately 500 foods from the USDA Database for the Choline Content of Common Foods (USDA, 2004). Values for other choline metabolites included in the choline database, as well as some values for food items where we are analyzing additional samples, were not incorporated into SR.
- Added about 50 traditional or subsistence foods collected for development of the American Indian/Alaska Native Foods Database. These foods were hunted, harvested, fished, and prepared by tribal members. Data developed for this project are shared with the tribes that provided the samples. In many cases, the wild form of a plant or animal has a very different nutrient profile than its domesticated counterpart (e.g., berries, fish). In almost all cases, these data are generated from a limited sampling. Therefore, they are subject to revision as additional samples are analyzed.
- A study was conducted to determine the nutrient composition of cured ham products. Nationally representative samples of various ham products were obtained from 12 locations nation-wide, prepared (unheated or heated according to package directions), and analyzed for nutrient content. As a result of this study, 74 new cured ham items have been added to SR. In addition to cured natural hams, products for enhanced categories are included. Ham products are defined as follows: “ham” contains at least 20.5% protein in the lean area with no water added; “ham with natural juices” contains at least 18.5% protein with a small addition of water when cured; “ham and water added” contains at least 17% protein with no more than 10% added solution; “ham and water product” contains less than 17% protein and contains water, but labeling must indicate percentage of "added ingredients."
- In addition to the foods mentioned above, a number of new foods were added to the database, including 30 breakfast cereals, 29 fast foods (breakfast items, hamburger and chicken sandwiches) and 26 sweets (desserts, frostings and candies) using data submitted by the food industry or generated by USDA through NFNAP. A complete list of the added food items is in the ADD\_FOOD file (p. 31). In addition, many nutrient values in all food groups have been updated with new data from NFNAP, food companies, and the scientific literature. These can be found in the CHG\_NUTR file (p. 31).
- A number of brand name items in Baked Products, Breakfast Cereals, and Meals, Entrees, and Sidedishes were deleted because products were no longer on the market or nutritional information does not reflect current market data.

## **Data Files**

The data files for SR19 are in ASCII format. A description of each field in these files and the relationships between each begins on p. 20. We have also made a Microsoft Access 2000 database available. This database contains all the SR19 files and relationships, with a few sample queries and reports. We have also included an abbreviated file (p. 28), with fewer nutrients but all the food items. A Microsoft Excel 2000 spreadsheet of this file is also provided.

## **Reports**

The data in SR19 are also available as reports in two different presentations. The first presents items in SR19 as page images containing all the data for each food. These data are separated into files by food groups. The second presentation contains selected foods and nutrients in SR19. These reports are sorted either by food description or in descending order by nutrient content in terms of common household measures. The food items and weights in these reports are adapted from those in the “U.S. Department of Agriculture Home and Garden Bulletin 72, Nutritive Value of Foods” (Gebhardt and Thomas, 2002).

The Adobe Reader is needed to see these files. There is a link from the NDL website to Adobe's website where it can be downloaded at no charge.

## **File Content**

As mentioned earlier, the database consists of several separate data files. The sections below provide details about the information in each. More extensive details on many specific foods are available in the printed “Agriculture Handbook 8” sections (U.S. Department of Agriculture, 1976-92).

The four principal files are the Food Description file, Nutrient Data file, Gram Weight file, and Footnote file. The six support files are the Nutrient Definition file, Food Group Description file, Source Code file, Data Derivation Code Description file, Sources of Data file, and Sources of Data Link file.

### **Food Description File**

This file includes descriptive information about the food items. For more details on the Food Description file, see “Food Description File Formats” (p. 21). A full description and a short description (containing abbreviations) are provided. Abbreviations used in creating short descriptions are given in Appendix A. In creating the short description, we did not abbreviate the first word in the long description. In addition, if the long description was 25 characters or less, the short description contains no abbreviations. Abbreviations used elsewhere are given in Appendix B. Brand names used in food descriptions are in upper case. Scientific names, common names, manufacturers' names, amounts of refuse, and refuse descriptions are provided where appropriate. The food group to which the food item belongs is also indicated. A code is also provided indicating if an item is used in the USDA Food and Nutrient Database for Dietary Studies (FNDDS). The factors used to calculate protein from nitrogen are included, as well as those used to calculate calories. There are no factors for items prepared using the recipe program of the NDBS or for items where the manufacturer calculates protein and calories.

The refuse and refuse description fields contain amounts and descriptions of inedible material (for example, seeds, bone, and skin) for applicable foods. These amounts are expressed as a percentage of the total weight of the item as purchased, and they are used to compute the weight of the edible portion. Refuse data were obtained from USDA-sponsored contracts and U.S. Department of Agriculture Handbooks 102 and 456 (Adams, 1975; Matthews and Garrison,

1975). To calculate “amount of nutrient in edible portion of 1 pound as purchased,” use the following formula:

$$Y = V*4.536*[(100-R)/100]$$

where

Y = nutrient value per 1 pound as purchased,

V = nutrient value per 100 g (Nutr\_Val in the Nutrient Data file), and

R = percent refuse (Refuse in the Food Description file).

For meat cuts containing bone and connective tissue, the amount of connective tissue is included in the value given for bone. Separable fat is not shown as refuse if the meat is described as separable lean and fat. Separable fat generally refers to seam fat and intramuscular fat. Separable lean refers to muscle tissue that can be readily separated from the intact cut; it includes any fat striations (marbling) within the muscle. For boneless cuts, the refuse values apply to connective tissue or connective tissue plus separable fat. The percentage yield of cooked, edible meat from 1 pound of raw meat with refuse can be determined by using the following formula:

$$Y = (W_c/453.6)*100$$

where

Y = nutrient value per 1 pound as purchased, and

W<sub>c</sub> = weight of cooked, edible meat.

## **Nutrients**

The Nutrient file contains nutrient values per 100 g, edible portion, along with fields to further describe the mean value. Nutrient values have been rounded to the number of decimal places for each nutrient as specified in the Nutrient Definition file (p. 24). We provide the following statistical attributes to better describe the data:

- Nutrient value – the mean of the data values for a specific parameter.
- Number of data points – the number of data points used to estimate the mean.
- Standard error – the standard error of the mean; a measure of variability of the mean value.
- Number of studies—the number of analytical studies used to generate a mean. A study is a discrete research project on the analysis of foods. A study can be the analysis of one nutrient in one food, one nutrient in many foods, or many nutrients in many foods.
- Minimum value—the smallest observed value in a range of values.
- Maximum value—the largest observed value in a range of values.
- Degrees of freedom—the number of data values that are free to vary after we have placed certain restrictions on the estimates; used in probability calculations.
- Lower- and upper-error bounds—represent a range of values within which the population mean is expected to fall, given a pre-specified confidence level. For SR19 and related releases, the confidence level is 95 percent.
- Statistical comments—give additional details about certain assumptions made during

statistical calculations. The definition of each comment is given in the discussion of the Nutrient Value file under “File Formats” (p. 23).

Other fields provide information on how the values are generated, as follows:

- Derivation code—gives more information about how a value was calculated or imputed. Procedures for imputing nutrient values were described by Schakel et al. (1997).
- Reference NDB number—indicates the NDB number of the food item that was used to impute a nutrient value for another food. This field is only populated for items added or updated since SR14.
- Added nutrient marker—a “Y” indicates that a mineral or vitamin was added for enrichment or fortification. This field is populated for ready-to-eat breakfast cereals and many brand name hot cereals in food group 8. In future releases, this field will be populated for other food groups.
- Confidence code—indicates the relative quality of the data. This code is derived using the data quality criteria first described by Mangels et al. (1993). These criteria have been expanded and enhanced for the new NDBS (Holden et al., 2002). This field is included as a placeholder for future releases.

For more details on the Nutrient Data file, see “Nutrient Data File Formats” (p. 23). Nutrient values indicate the total amount of the nutrient present in the edible portion of the food, including any nutrients added in processing. Table 1 gives an idea of the comprehensiveness of the database by listing for each nutrient the number of items that contain data.

When nutrient data for prepared or cooked products were unavailable or incomplete, nutrient values were calculated from comparable raw items or by recipe. When values are calculated in a recipe or from the raw item, appropriate nutrient retention (USDA, 2003) and yield factors (Matthews and Garrison, 1975) are applied. To obtain the content of nutrient per 100 g of cooked food, the nutrient content per 100 g of raw food is multiplied by the nutrient retention factor and, when appropriate, adjustments are made for fat and moisture gains and losses.

Nutrient retention factors are based on data from USDA research contracts, recent research reported in the literature, and USDA publications. Most retention factors were calculated by the True Retention Method (%TR) (Murphy et al., 1975). This method, as shown below, accounts for the loss of solids from foods that occurs during preparation and cooking.

$$\%TR = (N_c * G_c) / (N_r * G_r) * 100$$

where

$N_c$  = nutrient content per g of cooked food,  
 $G_c$  = g of cooked food,  
 $N_r$  = nutrient content per g of raw food, and  
 $G_r$  = g of food before cooking.



In general, levels of fortified nutrients are the values calculated by the manufacturer or NDL food specialists, based on the Nutrition Labeling and Education Act (NLEA) label declaration of % Daily Value (DV) (CFR, Title 21, Pt. 101) (U.S. Food and Drug Administration–Department of Health and Human Services, 2004). Such values represent the minimum nutrient level one can expect in the product. If analytical values were used to estimate levels of added nutrients, a number is present in the sample count field for these nutrients.

**Table 1.—Number of Foods in Database (n = 7,293)**

Nutr. No.	Nutrient	Number of foods	Nutr. No.	Nutrient	Number of foods
255	Water* †	7289	317	Selenium* †	6020
208	Energy* †	7293	313	Fluoride, F	509
203	Protein* †	7293	318	Vitamin A (IU)* †	6994
204	Total lipid (fat)* †	7293	320	Vitamin A (RAE)* †	6280
207	Ash* †	7286	319	Retinol* †	6199
205	Carbohydrate, by difference* †	7293	321	β-carotene* †	4104
291	Total dietary fiber* †	6584	322	α-carotene* †	3993
209	Starch	472	334	β-cryptoxanthin* †	3981
269	Total sugars* †	4767	337	Lycopene* †	3950
210	Sucrose	854	338	Lutein+zeaxanthin* †	3894
211	Glucose	861	324	Vitamin D	510
212	Fructose	852	323	α-tocopherol (vitamin E)* †	4183
213	Lactose	814	573	Added vitamin E*	3204
214	Maltose	801	341	β-tocopherol	1003
287	Galactose	687	342	γ-tocopherol	1014
301	Calcium* †	7166	343	δ-tocopherol	1007
303	Iron* †	7192	430	Vitamin K* †	3808
304	Magnesium* †	6790	401	Vitamin C, total ascorbic acid* †	6969
305	Phosphorus* †	6894	404	Thiamin* †	6857
306	Potassium* †	6951	405	Riboflavin* †	6862
307	Sodium* †	7211	406	Niacin* †	6855
309	Zinc* †	6790	410	Pantothenic acid†	6066
312	Copper*	6702	415	Vitamin B <sub>6</sub> * †	6631
315	Manganese†	5894			

\*Nutrients included in the USDA Food and Nutrient Database for Dietary Studies (FNDDS).

† Nutrients included in the Abbreviated file (p. 28)

**Table 1.—Number of Foods in Database (n = 7,293)—(continued)**

417	Folate, total* †	6610	687	17:1	715
431	Folic acid* †	6312	617	18:1 undifferentiated*	6323
432	Food folate* †	6419	674	18:1 c	277
435	Folate (DFE)* †	6304	663	18:1 t	289
421	Choline, total	478	628	20:1*	5323
454	Betaine	477	630	22:1 undifferentiated*	4569
418	Vitamin B <sub>12</sub> * †	6590	676	22:1 c	19
578	Added vitamin B <sub>12</sub> *	3267	664	22:1 t	6
601	Cholesterol* †	7039	671	24:1 c	186
636	Phytosterols	592	646	Total polyunsaturated fatty acids* †	6727
641	β-Sitosterol	68			
638	Stigmasterol	67	618	18:2 undifferentiated*	6341
639	Campesterol	67	675	18:2 n-6 c,c	252
221	Alcohol*	4251	666	18:2 i	78
262	Caffeine*	3923	669	18:2 t,t	141
263	Theobromine*	3902	665	18:2 t not further defined	10
606	Total saturated fatty acids* †	6981	670	18:2 CLAs	35
607	4:0*	4352	619	18:3 undifferentiated*	6222
608	6:0*	4377	851	18:3 n-3 c,c,c	456
609	8:0*	4643	685	18:3 n-6 c,c,c	430
610	10:0*	5323	856	18:3 i	32
611	12:0*	5647	627	18:4*	4354
696	13:0	203	672	20:2 n-6 c,c	751
612	14:0*	6069	689	20:3 undifferentiated	774
652	15:0	969	852	20:3 n-3	5
613	16:0*	6300	853	20:3 n-6	14
653	17:0	999	620	20:4 undifferentiated*	5341
614	18:0*	6286	855	20:4 n-6	9
615	20:0	1117	629	20:5 n-3*	4544
624	22:0	949	857	21:5	51
654	24:0	475	858	22:4	53
645	Total monounsaturated fatty acids* †	6720	631	22:5 n-3*	4488
			621	22:6 n-3*	4542
625	14:1	1051	605	Fatty acids, total <i>trans</i>	823
627	15:1	691	693	Fatty acids, total <i>trans</i> -monoenoic	267
697	16:1 undifferentiated*	6029			
673	16:1 c	88	695	Fatty acids, total <i>trans</i> -polyenoic	212
662	16:1 t	32			

\*Nutrients included in the USDA Food and Nutrient Database for Dietary Studies (FNDDS).

† Nutrients included in the Abbreviated file (p. 28)

**Table 1.—Number of Foods in Database (n = 7,293)—(continued)**

Nutr. No.	Nutrient	Number of foods	Nutr. No.	Nutrient	Number of foods
501	Tryptophan	4670	511	Arginine	4705
502	Threonine	4715	512	Histidine	4710
503	Isoleucine	4718	513	Alanine	4657
504	Leucine	4719	514	Aspartic acid	4660
505	Lysine	4733	515	Glutamic acid	4661
506	Methionine	4730	516	Glycine	4658
507	Cystine	4659	517	Proline	4646
508	Phenylalanine	4716	518	Serine	4656
509	Tyrosine	4684	521	Hydroxyproline	732
510	Valine	4718			

\*Nutrients included in the USDA Food and Nutrient Database for Dietary Studies (FNDDS).

† Nutrients included in the Abbreviated file (p. 28)

**Proximates.** The term proximate components refers to those macronutrients that include water (moisture), protein, total lipid (fat), total carbohydrate, and ash.

**Protein.** The values for protein were calculated from the level of total nitrogen (N) in the food, using the conversion factors recommended by Jones (1941). The specific factor applied to each food item is provided in the N\_Factor field in the Food Description file. The general factor of 6.25 is used to calculate protein in items that do not have a specific factor. There is no factor for items prepared using the recipe program of the NDBS or for items where the manufacturer calculates protein.

Protein values for chocolate, cocoa, coffee, mushrooms, and yeast were adjusted for nonprotein nitrogenous material. The adjusted protein conversion factors used to calculate protein for these items are as follows:

chocolate and cocoa	4.74
coffee	5.3
mushrooms	4.38
yeast	5.7

When these items were used as ingredients, such as chocolate in chocolate milk or yeast in bread, only their protein nitrogen content was used to determine their contribution to the calculated protein and amino acid content of the food. Protein calculated from total nitrogen, which may contain nonprotein nitrogen, was used in determining carbohydrate by difference. This unadjusted protein value is not given in the Nutrient Data file for SR19; rather, it is given as a footnote in printed sections of “Agriculture Handbook 8.”

For soybeans, nitrogen values were multiplied by a factor of 5.71 (Jones, 1941) to calculate protein. The soybean industry, however, uses 6.25 to calculate protein. The protein content of soy flours, soy meals, soy protein concentrates, and soy protein isolates is expressed both ways.

The item calculated using the 6.25 factor is identified as “crude protein basis.”

Total Lipid. The total lipid (fat) content of most foods was determined by gravimetric methods, including extraction methods such as those that use ether or a mixed solvent system of chloroform and methanol, or by acid hydrolysis. Total lipid determined by extraction is reported as Nutrient No. 204. It is sometimes referred to as “crude fat” and includes the weight of all lipid components, including glycerol, soluble in the solvent system. Nutrient No. 204 may not be identical to the fat level declared on food labels under the NLEA, where fat is expressed as the amount of triglyceride that would produce the analytically determined amount of lipid fatty acids and does not include other lipid components not soluble in the solvent system. The term “NLEA fat” is commonly referred to as “total fatty acids expressed as triglycerides.”

Carbohydrate. Carbohydrate, when present, was determined as the difference between 100 and the sum of the percentages of water, protein, total lipid (fat), ash, and, when present, alcohol. Total carbohydrate values include total dietary fiber. Carbohydrate in beer and wine was determined by methods 979.06 (27.1.21) and 985.10 (28.1.18) of AOAC International (AOAC 2003), respectively. Total dietary fiber content was determined by enzymatic-gravimetric methods 985.29 and 991.43 of the AOAC (2003). Total sugars is the term used for the sum of the individual monosaccharides (galactose, glucose, and fructose) and disaccharides (sucrose, lactose, and maltose). Analytical data for individual sugars were determined using AOAC methods (2003), either high-performance liquid chromatography (HPLC) or gas-liquid chromatography (GLC). When analytical data for total sugars were unavailable for items in the FNDDS, values were imputed or obtained from manufacturers and trade associations. Starch was analyzed using the AOAC method 966.11 (2003). Because the analyses of total dietary fiber, total sugars, and starch are performed separately and reflect the analytical variability inherent to the measurement process, the sum of these carbohydrate fractions may not equal the carbohydrate-by-difference value.

Food Energy. Food energy is expressed in kilocalories (kcal) and kilojoules (kJ). One kcal equals 4.184 kJ. The data are for physiological energy, which is the energy value remaining after losses from digestion and metabolism, are deducted from gross energy. Calorie values, with the exception of multi-ingredient processed foods, are based on the Atwater system for determining energy values. Derivation of the Atwater calorie factors is discussed in “Agriculture Handbook 74” (Merrill and Watt, 1973). For multi-ingredient processed foods, calorie values (source codes 8 or 9; for more information on source codes, see p. 25) generally reflect industry practices (as permitted by NLEA) of calculating calories from 4-4-9 calories per gram of protein, carbohydrate, and fat, respectively, or from 4-4-9 calories per gram of protein, carbohydrate minus insoluble fiber, and fat. The latter method is often used for high-fiber foods.

Calorie factors for protein, fat, and carbohydrates are included in the Food Description file. For foods containing alcohol, we used a factor of 6.93 to calculate calories per gram of alcohol. No calorie factors are given for items prepared using the recipe program of the NDBS. Instead, total calories for these items equal the sums of the calories contributed by each ingredient after adjustment for changes in yield, as appropriate. For multi-ingredient processed foods, if the calories calculated by the manufacturer are reported, no calorie factors are given.

Calorie factors for fructose and sorbitol, not available in the Atwater system, were derived from the work of Livesay and Marinos (1988). Calorie factors for coffee and tea were estimated from those for seeds and vegetables, respectively.

**Minerals.** Minerals included in the database are calcium, iron, magnesium, phosphorus, potassium, sodium, zinc, copper, manganese, selenium, and fluoride. Levels of minerals for most foods were determined by methods of the AOAC (2003). Calcium, iron, magnesium, phosphorus, sodium, potassium, zinc, copper, and manganese were usually determined by atomic absorption (AOAC 985.35) and inductively coupled plasma emission spectrophotometry (AOAC 984.27)

Analytical data for selenium were published earlier by USDA (1992) and were determined by the modified selenium hydride and fluorometric methods. Selenium values from the NFNAP were determined by either the modified selenium hydride (AOAC 986.15) or stable isotope dilution gas chromatography-mass spectrometry (Reamer and Veillon, 1981) methods. The selenium content of plants, in particular cereal grains, is strongly influenced by the quantity of biologically available selenium in the soil in which the plants grow, that is, by their geographical origin (Kubota and Allaway, 1972). The values given are national averages and should be used with caution when levels of selenium in locally grown foods are of interest or concern.

Values for fluoride, previously released in the USDA National Fluoride Database of Selected Beverages and Foods, Release 2 (USDA, 2005) have been incorporated into SR19, but other analyzed values, including regional values, are not included in SR. Samples were analyzed using a fluoride ion-specific electrode direct read method (VanWinkle, 1995) for clear liquids and a micro-diffusion method (VanWinkle, 1995) for other food samples.

**Vitamins.** Vitamins included in the database are ascorbic acid (vitamin C), thiamin, riboflavin, niacin, pantothenic acid, vitamin B<sub>6</sub>, vitamin B<sub>12</sub>, folate, total choline and betaine, vitamin A, vitamin E ( $\alpha$ -tocopherol), vitamin K (phylloquinone), and vitamin D.

Ascorbic acid. In the current database system, all data for ascorbic acid are listed under Nutrient No. 401, total ascorbic acid, determined by the fluorometric method (AOAC 967.22). Older values which have not been updated are primarily reduced ascorbic acid and were determined by the dichloroindophenol method (AOAC 967.21)

Thiamin, Riboflavin, and Niacin. Thiamin was determined chemically by the fluorometric method (AOAC 942.23). Fluorometric (AOAC 970.65) or microbiological (AOAC 940.33) methods were used to measure riboflavin. Niacin was determined by microbiological methods (AOAC 944.13). The values for niacin are for preformed niacin only and do not include the niacin contributed by tryptophan, a niacin precursor. The term “niacin equivalent” applies to the potential niacin value, that is, to the sum of the preformed niacin and the amount that could be derived from tryptophan. To estimate the amounts of niacin available from foods, the mean value of 60 mg tryptophan is considered equivalent to 1 mg niacin (IOM, 1998).

$$\text{mg Niacin equivalents} = \text{mg niacin} + (\text{mg tryptophan} / 60)$$

Pantothenic acid, Vitamins B<sub>6</sub>, and B<sub>12</sub>. Pantothenic acid was determined microbiologically (AOAC 945.74 or 992.07). Vitamin B<sub>6</sub> was determined by microbiological methods (AOAC 961.15) and B<sub>12</sub> was also determined by microbiological methods (AOAC 952.20). Vitamin B<sub>12</sub> is found in foods of animal origin or those containing some ingredient of animal origin, for example, cake that contains eggs or milk. For foods that contain only plant products, the value for vitamin B<sub>12</sub> is assumed to be zero. Some reports contain values for vitamin B<sub>12</sub> in certain fermented foods (beer, soy sauce, and miso). It is believed that this B<sub>12</sub> is synthesized not by the microorganisms responsible for the fermentation of the food, but rather by other contaminating microorganisms. Therefore, one should not consider these foods to be a consistent source of vitamin B<sub>12</sub> (Liem et al., 1977).

The Dietary Reference Intakes (DRI) for vitamin B<sub>12</sub> recommend that people older than 50 years meet their Recommended Dietary Allowances (RDA) mainly by consuming foods fortified with vitamin B<sub>12</sub> or a vitamin B<sub>12</sub>-containing supplement (IOM, 1998). Since vitamin B<sub>12</sub>, added as a fortificant, may provide a significant source of the vitamin in the diet, a nutrient number (#578) for “added vitamin B<sub>12</sub>” has been added to the database. In this release, there are about 230 foods fortified with vitamin B<sub>12</sub>. The vast majority are breakfast cereals, infant formulas, and plant-based meat substitutes. For these foods, the value for total vitamin B<sub>12</sub> was used for “added vitamin B<sub>12</sub>.” Only a few cereals containing a milk ingredient would contain any intrinsic vitamin B<sub>12</sub>. Milk-based infant formulas would contain intrinsic vitamin B<sub>12</sub>. However, infants are not the population of concern for intake of fortified vitamin B<sub>12</sub>. Plant-based meat substitutes would not contain intrinsic vitamin B<sub>12</sub>.

Folate. Values are reported for folic acid, food folate, and total folate reported as µg of dietary folate equivalents (DFEs). These varied folate forms are included as described in the DRI report issued by the Institute of Medicine of the National Academies (IOM, 1998). RDAs for folate are expressed in DFEs, which take into account the greater bioavailability of synthetic folic acid compared with naturally occurring food folate.

To calculate DFEs for any single food, it is necessary to have separate values for naturally occurring food folate and added synthetic folic acid in that item.

$$\mu\text{g DFE} = \mu\text{g food folate} + (1.7 * \mu\text{g folic acid})$$

Folate values for foods analyzed through NFNAP are generated using the trienzyme microbiological procedure (Martin et al., 1990). Microbiological methods measure total folate; for enriched foods, folic acid and food folate are not distinguished from each other. Therefore, to be able to calculate DFE, multi-ingredient enriched foods are analyzed by an additional microbiological procedure without enzymes to estimate the amount of added folic acid (Chun et al., 2006). Food folate is then calculated by difference.

The addition of folic acid to enriched cereal-grain products subject to standards of identity began in the United States on January 1, 1998 (CFR, Title 21, Pts. 136–137). These products include flour, cornmeal and grits, farina, rice, macaroni, noodles, bread, rolls, and buns. Folic acid may continue to be added (with some restrictions on amounts) to breakfast cereals, infant formulas, medical foods, food for special dietary use, and meal replacement products.

For unenriched foods, food folate would be equivalent to total folate since folic acid (pteroylmonoglutamic acid) occurs rarely in foods. Therefore, the same value with its number of data points and standard error, if present, is used for total folate and food folate. The folic acid value is assumed to be zero.

For enriched cereal-grain products with standards of identity (flour, cornmeal and grits, farina, rice, macaroni, noodles, bread, rolls, and buns), the folic acid value is calculated by subtracting the analytical folate value before fortification from the analytical value for the fortified product.

Enriched ready-to-eat (RTE) cereals have generally included folic acid fortification for over 25 years. Therefore, food folate values (before fortification) were not readily available for these products. Food folate was estimated by means of the NDBS formulation program for a variety of high-consumption cereals. Mean folate values were calculated for categories of RTE cereals based on grain content. Added folic acid was then calculated by subtracting estimated food folate from the total folate content. Generally, food folate values represent a small proportion of the total folate in the fortified products.

Choline. Beginning with SR19, total choline and betaine values from the USDA Database for the Choline Content of Common Foods (USDA, 2004) were incorporated into SR. Values for the individual metabolites were not added to SR, but are available in the USDA Database for the Choline Content of Common Foods.

Choline compounds were extracted and partitioned into organic and aqueous phases using methanol and chloroform and analyzed directly by liquid chromatography - electrospray ionization -isotope dilution mass spectrometry (LC-ESI-IDMS) (Koc et al., 2002). Samples were analyzed for betaine and these choline-contributing compounds: free choline (Cho), glycerophosphocholine (GPC), phosphocholine (Pcho), phosphatidylcholine (Ptdcho), and sphingomyelin (SM).

Because there are metabolic pathways for the interconversion of Cho, GPC, Pcho, PtdCho, and SM (Zeisel et al., 1994), total choline content is calculated as the sum of these choline-contributing metabolites. Betaine values are not included in the calculation of total choline since the conversion of choline to betaine is irreversible (Zeisel et al., 2003).

Vitamin A. Beginning with SR15 (2002), in addition to the international units (IUs) that have been reported in the past, we reported values for vitamin A in  $\mu\text{g}$  of retinol activity equivalents (RAEs) and  $\mu\text{g}$  of retinol. Values in  $\mu\text{g}$  of retinol equivalents (REs) were dropped from the database.

This change responds to new reference values for vitamin A issued by the Institute of Medicine of the National Academies (IOM, 2001). Along with the new DRIs, the panel recommended changing the factors used for calculating vitamin A activity from the individual provitamin A carotenoids and introduced RAE as a new unit for expressing vitamin A activity. One  $\mu\text{g}$  RAE is equivalent to 1  $\mu\text{g}$  of all-*trans*-retinol, 12  $\mu\text{g}$  of all-*trans*- $\beta$ -carotene, or 24  $\mu\text{g}$  of other provitamin A carotenoids. The RAE conversion factors are based on recent studies that show that the

conversion of provitamin A carotenoids to retinol is only half as great as previously thought.

Vitamin A in IU will continue to be reported because it is still the unit used for nutrition labeling. One IU is equivalent to 0.3 µg retinol, 0.6 µg β-carotene, or 1.2 µg other provitamin-A carotenoids (NAS/NRC, 1989).

Individual carotenoids, β-carotene, α-carotene, β-cryptoxanthin, lycopene, and lutein+zeaxanthin are reported. The analytical data are from NFNAP, generated using HPLC methodology (AOAC 941.15) and from the scientific literature. Most analytical systems do not separate lutein and zeaxanthin, so these carotenoids are shown combined. These values supersede those in Holden et al., 1999. Vitamin A values in IU and RAE were calculated from the individual carotenoids (β-carotene, α-carotene, and β-cryptoxanthin) using the appropriate factors. For food items used in the FNDDS, carotenoid values were imputed if analytical data were not available. For many of these items data were only available for vitamin A in IU. However, the variability in carotenoid levels due to cultivar, season, growing area, etc., as well as rounding within the NDBS, increases the difficulty in matching the calculated vitamin A values from imputed individual carotenoids to the existing IU values. As a result, the vitamin A IU value should agree within ±15 IU of the value calculated from individual carotenoids.

When individual carotenoids are not reported for plant foods (such as fruits, vegetables, legumes, nuts, cereal grains, and spices and herbs), µg RAE were calculated by dividing the IU value by 20. In foods of animal origin, such as eggs, beef, pork, poultry, lamb, veal, game, and fish (except for some organ meats and dairy), all of the vitamin A activity is contributed by retinol. For these foods, when analytical data were not available, µg RAE and µg of retinol were calculated by dividing the IU value by 3.33.

In foods that contain both retinol and provitamin A carotenoids, the amount of each of these components must be known to calculate RAE. Most of the vitamin A data in the database were received as IU. Therefore, the amounts of the provitamin A carotenoids and retinol had to be estimated based on the amount of retinol and provitamin A carotenoids in the ingredients. Once the components had been estimated, µg RAE were calculated as (IU from carotenoids/20) + (IU from retinol/3.33). µg of retinol were calculated as IU from retinol/3.33.

Vitamin E. Vitamin E activity for the RDA as defined by the DRI report (IOM, 2000) is now limited to the naturally occurring form and three synthetic forms of α-tocopherol. For this reason, α-tocopherol equivalents, which included vitamin E activity from α-, β-, γ-, and δ-tocopherols and α-, β-, and γ-tocotrienols, were dropped from the database in SR16. Tocopherols were determined by gas-liquid chromatography (GLC) or high-performance liquid chromatography (HPLC) (Lee et al., 1999). For those items in FNDDS, values are presented for α-tocopherol. If analytical data were unavailable, values for α-tocopherol were imputed. When available, values are also presented for the other tocopherols.

In the DRI report for Vitamin E, different factors were used to calculate the milligram amount of α-tocopherol from IU of vitamin E (IOM, 2000). The factors vary depending upon the chemical form of α-tocopherol used to fortify the food where



mg of  $\alpha$ -tocopherol in food, fortified food, or multivitamin

= IU of the RRR- $\alpha$ -tocopherol compound  $\times$  0.67 and

= IU of the *all rac*- $\alpha$ -tocopherol compound  $\times$  0.45.

Before SR16-1, the conversion factor for RRR- $\alpha$ -tocopherol was used for all vitamin E fortified foods. New  $\alpha$ -tocopherol values have been calculated for breakfast cereals, most infant formulas and a few other foods that are fortified with vitamin E, where we have confirmed that all *rac*- $\alpha$ -tocopherol was the form added. For more information about vitamin E in breakfast cereals, see the article in the January 2004 issue of the *American Journal of Clinical Nutrition* by Leonard et al. (2004).

The basis of the vitamin E tolerable upper intake level (UL), another reference value in the DRI report, was established using all forms of supplemental  $\alpha$ -tocopherol (IOM, 2000).” A new nutrient number (#573) has been added to identify quantities of “added vitamin E.” In this release, there are 100 food items that have values for added vitamin E greater than 0. For the majority of these food items the form added is synthetic vitamin E (all *rac*- $\alpha$ -tocopherol). To relate intakes of supplemental  $\alpha$ -tocopherol to the UL, values for “added vitamin E” should be multiplied by 2 when the added form is synthetic vitamin E. Although the 2S-stereoisomers do not contribute to vitamin E activity for the RDA (IOM, 2000), they do contribute to the UL. Items that are fortified with RRR- $\alpha$ -tocopherol (natural vitamin E) are identified by a footnote and the added vitamin E can be used directly for contribution to the UL. The majority of foods that are fortified with vitamin E are infant formulas and breakfast cereals. For these foods, the value for total vitamin E was used for “added vitamin E”; the small amount of intrinsic vitamin E was not considered. In fortified peanut butter, the intrinsic vitamin E was calculated since it is a substantial amount.

Vitamin K. Much of the data for vitamin K were generated under NFNAP and supersede the values in the Provisional Table (PT-104) (Weihrauch and Chatra, 1994). Vitamin K is extracted with hexane, purified with solid phase extraction using silica columns, and quantitated using HPLC with chemical reduction and fluorescence detection. Losses are corrected using vitamin K<sub>1(25)</sub> as internal standard (Booth et al. 1994).

Vitamin D. Much of the data for vitamin D were published earlier in the Provisional Table (PT-108) (Weihrauch and Tamaki, 1991). Values for breakfast cereals were updated based on data received from the food industry; values for other food items were updated using data generated under NFNAP using liquid chromatography (AOAC 995.05 or 982.29). These new values supersede those in PT-108.

**Lipid Components.** Fatty acids are expressed as the actual quantity of fatty acid in g per 100 g of food and do not represent fatty acids as triglycerides. Historically, most fatty acid data were obtained as the percentage of fatty acid methyl esters and determined by GLC analyses (AOAC 996.06). These data were converted to g fatty acid per 100 g total lipid using lipid conversion factors and then to g fatty acid per 100 g edible portion of food using the total lipid content. Details of the derivation of lipid conversion factors were published by Weihrauch et al., 1977.

In the redesigned NDBS, fatty acid data may be imported in a variety of units and converted within the system. No conversions are required if data are received as g fatty acid per 100 g edible portion of food. Data received as fatty acid esters and as triglycerides are converted to fatty acids using Sheppard conversion factors. Sheppard conversion factors are based on the molecular weights of the specific fatty acid and its corresponding esters (butyl or methyl) and triglyceride (Sheppard, 1992). When fatty acid data are received as percentages of fatty acid methyl esters, methyl esters are converted to fatty acids using Sheppard conversion factors and then multiplied by total lipid (Nutrient No. 204) to give g fatty acid per 100 g edible portion of food. Occasionally, total lipid values are available from a variety of data sources, but individual fatty acids are available from fewer references. In those cases, it may be necessary to normalize the individual fatty acids to the mean fat value of the food item. In the case of normalized fatty acids, the sum of the individual fatty acids will equal the mean fat value multiplied by the Weihrauch (1977) lipid conversion factor for that food item. No statistics of variability are reported for normalized fatty acids.

Individual Fatty Acids. The basic format for describing individual fatty acids is that the number before the colon indicates the number of carbon atoms in the fatty acid chain, and the number after the colon indicates the number of double bonds. For unsaturated fatty acids, additional nutrient numbers have been added to accommodate the reporting of many specific positional and geometric isomers. Of the specific isomers, there are two basic classifications considered: omega double bond position and *cis/trans* configuration of double bonds.

Omega-3 and omega-6 isomers are denoted in shorthand nomenclature as n-3 and n-6. The n-number indicates the position of the first double bond from the methyl end of the carbon chain. The letter *c* or *t* indicates whether the bond is *cis* or *trans*. For polyunsaturated fatty acids, *cis* and *trans* configurations at successive double bonds may be indicated. For example, linoleic acid is an 18 carbon omega-6 fatty acid with 2 double bonds, both in *cis* configuration. When data are isomer specific, linoleic acid is described as 18:2 n-6 c,c. Other isomers of 18:2, for which new nutrient numbers have been assigned, include 18:2 c,t, 18:2 t,c, and 18:2 t,t; 18:2 t not further defined and 18:2 i. 18:2 i is not a single isomer but includes isomers other than 18:2 n-6 c,c with peaks that cannot be easily differentiated in the particular food item. Systematic and common names for fatty acids are given in Table 2.

Table 2 is provided for the convenience of users in attaching common names or systematic names to fatty acids in our database. Though individual fatty acids are more specific than in past releases, it is not possible to include every possible geometric and positional isomer. Where specific isomers exist for a fatty acid, the common name of the most typical isomer is listed for the undifferentiated fatty acid and an asterisk (\*) designates the specific isomer by that name. The most typical isomer for 18:1 is oleic. Thus, the specific isomer by that name, 18:1 c, is designated in Table 2 as oleic.

**Table 2.—Systematic And Common Names for Fatty Acids**

<b>Fatty acid</b>	<b>Systematic name</b>	<b>Common name of most typical isomer</b>	<b>Nutrient number</b>
Saturated fatty acids			
4:0	butanoic	butyric	607
6:0	hexanoic	caproic	608
8:0	octanoic	caprylic	609
10:0	decanoic	capric	610
12:0	dodecanoic	lauric	611
13:0	tridecanoic		696
14:0	tetradecanoic	myristic	612
15:0	pentadecanoic		652
16:0	hexadecanoic	palmitic	613
17:0	heptadecanoic	margaric	653
18:0	octadecanoic	stearic	614
20:0	eicosanoic	arachidic	615
22:0	docosanoic	behenic	624
24:0	tetracosanoic	lignoceric	654
Monounsaturated fatty acids			
14:1	tetradecenoic	myristoleic	625
15:1	pentadecenoic		697
16:1 undifferentiated	hexadecenoic	palmitoleic	626
16:1 c			673*
16:1 t			662
17:1	heptadecenoic		687
18:1 undifferentiated	octadecenoic	oleic	617
18:1 c			674*
18:1 t			663
20:1	eicosenoic	gadoleic	628
22:1 undifferentiated	docosenoic	erucic	630
22:1 c			676*
24:1 c	cis-tetracosenoic	nervonic	671
Polyunsaturated fatty acids			
18:2 undifferentiated	octadecadienoic	linoleic	618
18:2 n-6 c,c			675*
18:2 i			666
18:2 t,t			669
18:2 t not further defined			665

\* Designates the specific isomer associated with the common name; the typical isomer is listed for the undifferentiated fatty acid.

**Table 2.—Systematic And Common Names for Fatty Acids—(continued)**

<b>Fatty acid</b>	<b>Systematic name</b>	<b>Common name of most typical isomer</b>	<b>Nutrient number</b>
18:3	octadecatrienoic	linolenic	619
18:3 n-3 c,c,c		alpha-linolenic	851*
18:3 n-6 c,c,c		gamma-linolenic	685
18:4 undifferentiated	octadecatetraenoic	parinaric	627
20:2 n-6 c,c	eicosadienoic		672
20:3 undifferentiated	eicosatrienoic		689
20:3 n-3			852
20:3 n-6			853
20:4 undifferentiated	eicosatetraenoic	arachidonic	620
20:4 n-6			855
20:5 n-3	eicosapentaenoic (EPA)	timnodonic	629
21:5			857
22:4			858
22:5 n-3	docosapentaenoic (DPA)	clupanodonic	631
22:6 n-3	docosahexaenoic (DHA)		621

\* Designates the specific isomer associated with the common name; the typical isomer is listed for the undifferentiated fatty acid.

Fatty acid totals. Only a small portion of the fatty acid data received for release in SR19 contains specific positional and geometric isomers. Therefore, it has been necessary to maintain the usual nutrient numbers corresponding to fatty acids with no further differentiation other than carbon length and number of double bonds. To aid users of our data, specific isomers are always summed to provide a total value for the undifferentiated fatty acid. Thus, mean values for the specific isomers of 18:2 would be summed to provide a mean for 18:2 undifferentiated (Nutrient No. 618). Other fatty acid totals provided are (1) the sum of saturated, monounsaturated, and polyunsaturated fatty acids and (2) the sum of *trans*-monoenoic, the sum of *trans*-polyenoic, and the sum of all *trans* fatty acids.

Values for total saturated, monounsaturated, and polyunsaturated fatty acids may include individual fatty acids not reported; therefore, the sum of their values may exceed the sum of the individual fatty acids. In rare cases, the sum of the individual fatty acids may exceed the sum of the values given for the total saturated fatty acids (SFA), monounsaturated fatty acids (MUFA), and polyunsaturated fatty acids (PUFA). These differences are generally caused by rounding and may be relatively small.

For multi-ingredient processed brand-name foods, industry data were often available for fatty acid classes (SFA, MUFA, and PUFA) but were lacking for individual fatty acids. In these cases, individual fatty acids were calculated from the fatty acids of the individually listed ingredients and normalized to the total fat level. A best-fit approximation was made to fatty acid classes, but unavoidably, calculated sums of individual fatty acid totals did not always match industry data

for fatty acid classes. Zero values for individual fatty acids should be understood to mean that trace amounts may be present. When g fatty acids per 100 g of total lipid were converted to g fatty acids per 100 g of food, values of less than 0.0005 were rounded to 0.

Cholesterol. Cholesterol values were generated primarily by GLC procedures (AAOC 994.10). It is assumed that cholesterol is present only in foods of animal origin and foods containing at least one ingredient of animal origin (for example, cake that contains eggs). For mixtures containing ingredients derived from animal products, the cholesterol value may have been calculated from the value for those ingredients. For foods that contain only plant products, the value for cholesterol is assumed to be zero.

Plant sterols. Data on plant sterols (campesterol, stigmasterol, and  $\beta$ -sitosterol) were obtained by gas-chromatographic procedures (AOAC 967.18) and summed to calculate total phytosterols.

**Amino Acids**. Amino acid data for a class or species of food are aggregated to yield a set of values that serve as the pattern for calculating the amino acid profile of other similar foods. The amino acid values for the pattern are expressed on a per-gram-of-nitrogen basis. Amino acids are extracted in three groups—tryptophan, sulfur-containing amino acids (methionine and cystine), and all others. Tryptophan is determined by alkaline hydrolysis/HPLC (AOAC 988.15), methionine and cystine by performic oxidation/HPLC (AOAC 994.12) and all others by acid hydrolysis/HPLC (AOAC 982.30). The amino acid patterns and the total nitrogen content were used to calculate the levels of individual amino acids per 100 g of food, using the following formula:

$$AA_f = (AA_n * V_p) / N_f$$

where

$AA_f$  = amino acid content per 100 g of food,

$AA_n$  = amino acid content per g of nitrogen,

$V_p$  = protein content of food, and

$N_f$  = nitrogen factor.

In the past, the number of data points appeared only on the food item for which the amino acid pattern was developed, not on other foods that used the same pattern. It referred to the number of observations used in developing the amino acid pattern for that food. For foods processed in the new NDBS, the number of observations used in developing an amino acid pattern will be released only with the pattern. The amino acid profiles calculated from these patterns will show the number of data points to be zero.

If amino acid values are presented for an item with more than one protein-containing ingredient, the values may have been calculated on a per-gram-of-nitrogen basis from the amino acid patterns of the various protein-containing ingredients. Then the amino acid contents for an item on the 100-g basis were calculated as the sum of the amino acids in each protein-containing ingredient multiplied by total nitrogen in the item.

## **Weights and Measures**

Information is provided on household measures for food items (for example, 1 cup, 1 tablespoon, 1 fruit, 1 leg). Weights are given for edible material without refuse, that is, the weight of an apple without the core or stem, or a chicken leg without the bone, and so forth. The Weight file contains the gram weights and measure descriptions for each food item. This file can be used to calculate nutrient values for food portions from the values provided per 100 g of food. The following formula is used to calculate the nutrient content per household measure:

$$N = (V*W)/100$$

where

N = nutrient value per household measure,  
V = nutrient value per 100 g (Nutr\_Val in the Nutrient Data file), and  
W = g weight of portion (Gm\_Wgt in the Weight file).

The Weight file can be used to produce reports showing the household measure and nutrient values calculated for that portion. The weights were derived from published sources, industry files, studies conducted by USDA (Adams, 1975; Fulton et al., 1977), and the weights and measures used in the FNDDS (2004). Though we made a special effort to provide representative values, weights and measures obtained from different sources vary considerably for some foods. The format of this file is described on p. 26.

## **Footnotes**

Footnotes are provided for a few items where information about food description, weights and measures, or nutrient values could not be accommodated in existing fields. For example, if citric acid was added to a juice drink, this is indicated in the footnote. The format of this file is described on p. 27.

## **Sources of Data**

We first added this file (previously called References) with SR14 (2001). We have changed the name and some of the fields to reflect the fact that not all sources are journals or published literature, but also include the results of unpublished data from USDA-sponsored research and from research sponsored by others either separately or in collaboration with USDA. It contains data sources for the nutrient values and links to an identification number on each nutrient record. Since many of the data in this release were carried forward from SR13 (1999), nutrient-specific source documentation was not electronically available. As new data for these foods are generated and as additional documentation is entered into the new NDBS, data source information will increase in future releases. The format of this file is described on p. 27.

A file, the Sources of Data Link file, is provided to allow users to establish a relationship between the Sources of Data file and the Nutrient Data file. This lets the user identify specific sources of data for each nutrient value. For example, the user can use these files to identify

where NFNAP data is used in the database. The format of this file is described on p. 28.

### **Explanation of File Formats**

The data appear in two different organizational formats. One is a relational format of four principal and six support files making up the database. The relational format is complete and contains all food, nutrient, and related data. The other is a flat abbreviated file with all the food items, but fewer nutrients, and not all of the other related information. The abbreviated file does not include values for starch, fluoride, total choline and betaine, added vitamin E and added vitamin B<sub>12</sub>, alcohol, caffeine, theobromine, vitamin D, phytosterols, or individual amino acids, fatty acids, and sugars.

### **Relational Files**

The four principal and six support files of the relational database can be linked together in a variety of combinations to produce queries and generate reports (Figure 1). Table 3 shows the number of records in each file.

**Table 3.—Number of Records in Principal and Support Files**


<b>File name</b>	<b>Table name</b>	<b>Number of records</b>
<b>Principal files</b>		
Food Description	FOOD_DES	7,293
Nutrient Data	NUT_DATA	505,702
Weight	WEIGHT	13,064
Footnote	FOOTNOTE	247
<b>Support files</b>		
Food Group Description	FD_GROUP	24
Nutrient Definition	NUTR_DEF	140
Source Code	SRC_CD	10
Data Derivation Description	DERIV_CD	54
Sources of Data	DATA_SRC	425
Sources of Data Link	DATSRCLN	113,575

The relational files are in both ASCII format and a Microsoft Access 2000 database. Tables 4–13 describe the formats of these files. Information on the relationships that can be made among these files is also given. Fields that always contain data and fields that can be left blank or null are identified in the “blank” column; N indicates a field that is always filled; Y indicates a field that may be left blank (null) (Tables 4–13). An asterisk (\*) indicates that this is an index field for the file. Though the ASCII files are not indexed, the file descriptions show where indices were used to sort and manage records within the NDBS. When importing these files into a database management system, if files are to be indexed, it is important to use the indices listed here,

particularly with the Nutrient Data file, which uses two.

ASCII files are delimited. All fields are separated by carets (^) and text fields are surrounded by tildes (~). A double caret (^) or two carets and two tildes (~) appear when a field is null or blank. Format descriptions include the name of each field, its type [N = numeric with width and number of decimals (w.d) and A = alphanumeric], and maximum length. The actual length in the data files may be less and most likely will change in later releases. Values will be padded with zeroes when imported into various software packages, depending on the formats used.

**Figure 1.** Relationships among files in the USDA National Nutrient Database for Standard Reference



**Food Description File** (file name = FOOD\_DES). The Food Description file (Table 4) contains a long and short description and food group for 7,293 food items, along with common names, manufacturer name, scientific name, percentage and description of refuse, and factors used for calculating protein and calories, if applicable. Items used in the FNDDS are also identified.



- Links to the Food Group Description file by the FdGrp\_Cd field
- Links to the Nutrient Data file by the NDB\_No field
- Links to the Weight file by the NDB\_No field
- Links to the Footnote file by the NDB\_No field
- Links to Sources of Data file by the NDB\_No field through the Sources of Data Link file

**Table 4.—Food Description File Format**

<b>Field name</b>	<b>Type</b>	<b>Blank</b>	<b>Description</b>
NDB_No	A 5*	N	5-digit Nutrient Databank number that uniquely identifies a food item
FdGrp_Cd	A 4	N	4-digit code indicating food group to which a food item belongs
Long_Desc	A 200	N	200-character description of food item
Shrt_Desc	A 60	N	60-character abbreviated description of food item. Generated from the 200-character description using abbreviations in appendix A. If short description was longer than 60 characters, additional abbreviations were made.
ComName	A 100	Y	Other names commonly used to describe a food, including local or regional names for various foods, for example, “soda” or “pop” for “carbonated beverages”
ManufacName	A 50	Y	Indicates the company that manufactured the product, when appropriate
Survey	A 1	Y	Indicates if the food item is used in the USDA Food and Nutrient Database for Dietary Studies (FNDDS) and has a complete nutrient profile for a specified set of nutrients
Ref_desc	A 60	Y	Description of inedible parts of a food item (refuse), such as seeds or bone
Refuse	N 2	Y	Percentage of refuse
SciName	A 60	Y	Scientific name of the food item. Given for the least processed form of the food (usually raw), if applicable
N_Factor	N 4.2	Y	Factor for converting nitrogen to protein (see p. 8)
Pro_Factor	N 4.2	Y	Factor for calculating calories from protein (see p. 9)
Fat_Factor	N 4.2	Y	Factor for calculating calories from fat (see p. 9)
CHO_Factor	N 4.2	Y	Factor for calculating calories from carbohydrate (see p. 9)

\* Index field for the Food Description file.

**Food Group Description File** (file name = FD\_GROUP). This file (Table 5) is a support file to the Food Description file and contains a list of food groups used in SR19 and their descriptions.

- Links to the Food Description file by FdGrp\_Cd

**Table 5.—Food Group Description File Format**

Field name	Type	Blank	Description
FdGrp_Cd	A 4*	N	4-digit code identifying a food group. Only the first 2 digits are currently assigned. In the future, the last 2 digits may be used. Codes may not be consecutive.
FdGrp_Desc	A 60	N	Name of food group

\* Index field for the Food Group Description file.

**Nutrient Data File** (file name = NUT\_DATA). The Nutrient Data file (Table 6) contains the nutrient values and information about the values, including expanded statistical information.

- Links to the Food Description file by NDB\_No.
- Links to the Weight file by NDB\_No.
- Links to the Footnote file by NDB\_No and when applicable, Nutr\_No.
- Links to the Nutrient Definition file by Nutr\_No.
- Links to the Source Code file by Src\_Cd
- Links to the Derivation Code file by Deriv\_Cd

**Table 6.—Nutrient Data File Format**

Field name	Type	Blank	Description
NDB_No	A 5*	N	5-digit Nutrient Databank number
Nutr_No	A 3*	N	Unique 3-digit identifier code for a nutrient
Nutr_Val	N 10.3	N	Amount in 100 grams, edible portion †
Num_Data_Pts	N 5.0	N	Number of data points (previously called Sample_Ct)
Std_Error	N 8.3	Y	Standard error of the mean. Null if could not be calculated
Src_Cd	A 2	N	Code indicating type of data
Deriv_Cd	A 4	Y	Data Derivation Code giving specific information on how the value was determined
Ref_NDB_No	A 5	Y	NDB number of the item used to impute a missing value. Populated only for items added or updated starting with SR14

Add_Nutr_Mark	A 1	Y	Indicates a vitamin or mineral added for fortification or enrichment. This field is populated for ready-to-eat breakfast cereals and many brand name hot cereals in food group 8.
Num_Studies	N 2	Y	Number of studies
Min	N 10.3	Y	Minimum value
Max	N 10.3	Y	Maximum value
DF	N 2	Y	Degrees of Freedom
Low_EB	N 10.3	Y	Lower 95% error bound
Up_EB	N 10.3	Y	Upper 95% error bound
Stat_cmt	A 10	Y	Statistical comments. See definitions below.
CC	A 1	Y	Confidence Code indicating data quality, based on evaluation of sample plan, sample handling, analytical method, analytical quality control, and number of samples analyzed. Not included in this release, but is planned for future releases.

---

\* Index field for the Nutrient Data file.

† Nutrient values have been rounded to a specified number of decimal places for each nutrient. Number of decimal places is listed in the Nutrient Definition file.

Definitions of each Statistical Comment included in the Nutrient Data table follow:

1. The displayed summary statistics were computed from data containing some less-than values. Less-than, trace, and not-detected values were calculated.
2. The displayed degrees of freedom were computed using Satterthwaite's approximation (Korz and Johnson, 1988).
3. The procedure used to estimate the reliability of the generic mean requires that the data associated with each study be a simple random sample from all the products associated with the given data source (for example, manufacturer, variety, cultivar, and species).
4. For this nutrient, one or more data sources had only one observation. Therefore, the standard errors, degrees of freedom, and error bounds were computed from the between-group standard deviation of the weighted groups having only one observation.

**Nutrient Definition File** (file name = NUTR\_DEF). The Nutrient Definition file (Table 7) is the support file to the Nutrient Data file. It provides the 3-digit nutrient code, unit of measure, INFOODS tagname, and description.

- Links to the Nutrient Data file by Nutr\_No.

**Table 7.—Nutrient Definition File Format**

Field name	Type	Blank	Description
Nutr_No	A 3*	N	Unique 3-digit identifier code for a nutrient
Units	A 7	N	Units of measure (mg, g, µg, and so on.)
Tagname	A 20	Y	International Network of Food Data Systems (INFOODS) Tagnames.† A unique abbreviation for a nutrient/food component developed by INFOODS to aid in the interchange of data
NutrDesc	A 60	N	Name of nutrient/food component
Num_Dec	A 1	N	Number of decimal places to which a nutrient value is rounded
SR_Order	N 6	N	Used to sort nutrient records in the same order as various reports produced from SR

\* Index field for the Nutrient Definition file.

† INFOODS, 2006.

**Source Code File** (file name = SRC\_CD). The Source Code file (Table 8) contains codes indicating the type of data (analytical, calculated, assumed zero, and so on) in the Nutrient Data file. To improve the usability of the database and to provide values for the FNDDS, NDL staff imputed nutrient values for a number of proximate components, total dietary fiber, total sugar, and vitamin and mineral values.

- Links to the Nutrient Data file by Src\_Cd

**Table 8.—Source Code File Format**

Field name	Type	Blank	Description
Src_Cd	A 2*	N	2-digit code
SrcCd_Desc	A 60	N	Description of source code that identifies the type of nutrient data

\* Index field for the Source Code file.

**Data Derivation Code Description File** (file name = DERIV\_CD). This file (Table 9) is a support file for the Nutrient Data file and contains information on how the nutrient values were determined. The file contains the derivation codes and their descriptions.

- Links to the Nutrient Data file by Deriv\_Cd

**Table 9.—Data Derivation Code File Format**

Field name	Type	Blank	Description
Deriv_Cd	A 4*	N	Derivation Code
Deriv_Desc	A 120	N	Description of derivation code giving specific information on how the value was determined

\* Index field for the Data Derivation Code file.

For example, the data derivation code that indicates how  $\alpha$ -tocopherol (Nutrient No. 323) in Emu, fan fillet, raw (NDB. No. 05623) was calculated is BFSN. The breakdown of the code is as follows:

B = based on another form of the food or a similar food;  
 F = concentration adjustment used;  
 S = solids, the specific concentration adjustment used; and  
 N = retention factors not used

The Ref\_NDB\_No is 05621 Emu, ground, raw. This means that the analytical  $\alpha$ -tocopherol value in the total solids of emu, ground, raw is used to calculate the  $\alpha$ -tocopherol in the total solids of emu, fan fillet, raw.

$$N_t = (N_s * S_s) / S_t$$

where

$N_t$  = the nutrient content of the target item,

$N_s$  = the nutrient content of the source item,

For NDB No. 05621,  $\alpha$ -tocopherol = 0.24 mg/100 g

$S_s$  = the total solids content of the source item, and

For NDB No. 05621, solids = 25.38 g/100 g

$S_t$  = the total solids content of the target item.

For NDB No. 05623, solids = 27.13 g/100 g

So, using this formula for the above example:

$$N_t = (0.24 \times 25.38) / 27.13 = 0.22 \text{ mg/100 g } \alpha\text{-tocopherol in Emu, fan fillet, raw}$$

Only items that were imputed starting with SR14 (2001) will have both derivation codes and reference NDB numbers. Other items that were imputed outside the NDBS will have data derivation codes, but the Ref\_NDB\_No field will be blank.

**Weight File** (file name = WEIGHT). This file (Table 10) contains the weight in grams of a number of common measures for each food item.

- Links to Food Description file by NDB\_No.
- Links to Nutrient Data file by NDB\_No.

**Table 10.— Weight File Format**

Field name	Type	Blank	Description
NDB_No	A 5*	N	5-digit Nutrient Databank number
Seq	A 2*	N	Sequence number
Amount	N 5.3	N	Unit modifier (for example, 1 in “1 cup”)
Msre_Desc	A 80	N	Description (for example, cup, diced, and 1-inch pieces)
Gm_Wgt	N 7.1	N	Gram weight
Num_Data_Pts	N 3	Y	Number of data points
Std_Dev	N 7.3	Y	Standard deviation

\* Index field for the Weight file.

**Footnote File** (file name = FOOTNOTE). This file (Table 11) contains additional information about the food item, household weight, and nutrient value.

- Links to the Food Description file by NDB\_No.
- Links to the Nutrient Data file by NDB\_No and Nutr\_No.

**Table 11.—Footnote File Format**

Field name	Type	Blank	Description
NDB_No	A 5	N	5-digit Nutrient Databank number
Footnt_No	A 4	N	Sequence number. If a given footnote applies to more than one nutrient number, the same footnote number is used. As a result, this file cannot be indexed.
Footnt_Typ	A 1	N	Type of footnote. D = footnote adding information to the food description; M = footnote adding information to measure description; N = footnote providing additional information on a nutrient value. If the Footnt_typ = N, the Nutr_No will also be filled in
Nutr_No	A 3	Y	Unique 3-digit identifier code for a nutrient to which footnote applies
Footnt_Txt	A 200	N	Footnote text

**Sources of Data File** (file name = DATA\_SRC). This file (Table 12) provides a citation to the DataSrc\_ID in the Sources of Data Link file.

- Links to Nutrient Data file by NDB No. through the Sources of Data Link file

**Table 12.—Sources of Data File Format**

<b>Field name</b>	<b>Type</b>	<b>Blank</b>	<b>Description</b>
DataSrc_ID	A 6*	N	Unique number identifying the reference/source
Authors	A 255	Y	List of authors for a journal article or name of sponsoring organization for other documents
Title	A 255	N	Title of article or name of document, such as a report from a company or trade association
Year	A 4	Y	Year article or document was published
Journal	A 135	Y	Name of the journal in which the article was published
Vol_City	A 10	Y	Volume number for journal articles, books, or reports; city where sponsoring organization is located
Issue_State	A 5	Y	Issue number for journal article; State where the sponsoring organization is located
Start_Page	A 5	Y	Starting page number of article/document
End_Page	A 5	Y	Ending page number of article/document

\* Index field for the Sources of Data file.

**Sources of Data Link File** (file name = DATSRCLN). This file (Table 13) is used to link the Nutrient Data file with the Sources of Data table. It is needed to resolve the many-to-many relationship between the two tables.

- Links to the Nutrient Data file by NDB No. and Nutr\_No.
- Links to the Sources of Data file by DataSrc\_ID.

**Table 13.—Sources of Data Link File Format**

<b>Field name</b>	<b>Type</b>	<b>Blank</b>	<b>Description</b>
NDB_No	A 5*	N	5-digit Nutrient Databank number
Nutr_No	A 3*	N	Unique 3-digit identifier code for a nutrient
DataSrc_ID	A 6	N	Unique ID identifying the reference/source

\* Index field for the Sources of Data Link file.

### **Abbreviated File**

The Abbreviated file (file name = ABBREV) is available in ASCII format and is also provided in a Microsoft Excel spreadsheet. It contains all the food items, but fewer nutrients and other related information. The abbreviated file does not include values for starch, fluoride, total

choline and betaine, added vitamin E and added vitamin B<sub>12</sub>, alcohol, caffeine, theobromine, vitamin D, phytosterols, or individual amino acids, fatty acids, and sugars.

Table 14 lists all the nutrients included in the abbreviated file. Starting with SR16 (2003), we have added total sugars, folic acid, food folate, folate DFE, retinol, α-tocopherol, vitamin K, and individual carotenoids (α-carotene, β-carotene, β-cryptoxanthin, lycopene, and lutein+zeaxanthin) to the Abbreviated file. The ASCII file (Table 14) is in delimited format. Fields are separated by a caret (^). Text fields are surrounded by a tilde (~). Data refer to 100 g of the edible portion of the food item. Decimal points are included in the fields. Missing values are denoted by the null value of two consecutive carets (^) or two carets and two tildes (~). The file is sorted in ascending order by the NDB number. Two common measures are provided, which are the first two common measures in the Weight file for each NDB number. To obtain values per one of the common measures, multiply the value in the desired nutrient field by the value in the desired common measure field and divided by 100. For example, to calculate the amount of fat in 1 tablespoon of butter (NDB No. 01001),

$$V_H=(N*CM_2)/100$$

where:

V<sub>H</sub> = the nutrient content per the desired common measure

N = the nutrient content per 100 g

For NDB No. 01001, fat = 81.11 g/100 g

CM = grams of the common measure

For NDB No. 01001, 1 tablespoon = 9.4 g

So using this formula for the above example:

$$V_H = (81.11 * 9.4) / 100 = 7.52 \text{ g fat in 1 tablespoon of butter}$$

This file is a flat file and is provided for those users who do not need a relational database. It contains the information in one record per food item and is suitable for importing into a spreadsheet. We have imported the data file into a Microsoft Excel 2000 spreadsheet for users of that application. Users of other software applications can import either the Microsoft Excel 2000 spreadsheet or the ASCII files. It contains all the food items, but less descriptive information and fewer nutrients and weights than the larger relational files. If additional information is needed, this file can be linked to the other files by the NDB number.

**Table 14.—Abbreviated File Format**

Field name	Type	Description
NDB_No.	A 5*	5-digit Nutrient Databank number
Shrt_Desc	A 60	60-character abbreviated description of food item†
Water	N 10.2	Water (g/100 g)
Energ_Kcal	N 10	Food energy (kcal/100 g)
Protein	N 10.2	Protein (g/100 g)
Lipid_Tot	N 10.2	Total lipid (fat)(g/100 g)



<b>Field name</b>	<b>Type</b>	<b>Description</b>
Ash	N 10.2	Ash (g/100 g)
Carbohydr	N 10.2	Carbohydrate, by difference (g/100 g)
Fiber_TD	N 10.1	Total dietary fiber (g/100 g)
Sugar_Tot	N 10.2	Total sugars (g/100 g)
Calcium	N 10	Calcium (mg/100 g)
Iron	N 10.2	Iron (mg/100 g)
Magnesium	N 10	Magnesium (mg/100 g)
Phosphorus	N 10	Phosphorus (mg/100 g)
Potassium	N 10	Potassium (mg/100 g)
Sodium	N 10	Sodium (mg/100 g)
Zinc	N 10.2	Zinc (mg/100 g)
Copper	N 10.3	Copper (mg/100 g)
Manganese	N 10.3	Manganese (mg/100 g)
Selenium	N 10.1	Selenium ( $\mu\text{g}$ /100 g)
Vit_C	N 10.1	Vitamin C (mg/100 g)
Thiamin	N 10.3	Thiamin (mg/100 g)
Riboflavin	N 10.3	Riboflavin (mg/100 g)
Niacin	N 10.3	Niacin (mg/100 g)
Panto_acid	N 10.3	Pantothenic acid (mg/100 g)
Vit_B6	N 10.3	Vitamin B <sub>6</sub> (mg/100 g)
Folate_Tot	N 10	Folate, total ( $\mu\text{g}$ /100 g)
Folic_acid	N 10	Folic acid ( $\mu\text{g}$ /100 g)
Food_Folate	N 10	Food folate ( $\mu\text{g}$ /100 g)
Folate_DFE	N 10	Folate ( $\mu\text{g}$ dietary folate equivalents/100 g)
Vit_B12	N 10.2	Vitamin B <sub>12</sub> ( $\mu\text{g}$ /100 g)
Vit_A_IU	N 10	Vitamin A (IU/100 g)
Vit_A_RAE	N 10	Vitamin A ( $\mu\text{g}$ retinol activity equivalents/100g)
Retinol	N 10	Retinol ( $\mu\text{g}$ /100 g)
Vit_E	N 10.2	Vitamin E (alpha-tocopherol) (mg/100 g)
Vit_K	N 10.1	Vitamin K (phylloquinone) ( $\mu\text{g}$ /100 g)
Alpha_Carot	N 10	Alpha-carotene ( $\mu\text{g}$ /100 g)
Beta_Carot	N 10	Beta-carotene ( $\mu\text{g}$ /100 g)

<b>Field name</b>	<b>Type</b>	<b>Description</b>
Beta_Crypt	N 10	Beta-cryptoxanthin (µg/100 g)
Lycopene	N 10	Lycopene (µg/100 g)
Lut+Zea	N 10	Lutein+zeaxanthin (µg/100 g)
FA_Sat	N 10.3	Saturated fatty acid (g/100 g)
FA_Mono	N 10.3	Monounsaturated fatty acids (g/100 g)
FA_Poly	N 10.3	Polyunsaturated fatty acids (g/100 g)
Cholestrl	N 10.3	Cholesterol (mg/100 g)
GmWt_1	N 9.2	First household weight for this item from the Weight file‡
GmWt_Desc1	A 120	Description of household weight number 1
GmWt_2	N 9.2	Second household weight for this item from the Weight file‡
GmWt_Desc2	A 120	Description of household weight number 2
Refuse_Pct	N 2	Percent refuse§

\* Index field for the Abbreviated file.

† For a 200-character description and other descriptive information, link to the Food Description file.

‡ For the complete list and description of the measure, link to the Weight file.

§ For a description of refuse, link to the Food Description file.

### **Update Files**

These update files provide changes made between SR18 (2005) and SR19 (2006). Update files in ASCII are provided for those users who reformatted previous releases for their systems and wish to do their own updates. If you are using an earlier release, you will need to first obtain the update files for that release through SR18, update your database to SR18, then use the update files for SR19. These updates are available on NDL's website:

<http://www.ars.usda.gov/nutrientdata>. Added items are given in five files:

- ADD\_FOOD for descriptions of the new items,
- ADD\_NUTR for added nutrient data,
- ADD\_WGT for added weight and measure data,
- ADD\_FTNT for added footnotes,
- ADD\_NDEF for added nutrient definitions

These files are in the same formats as the Food Description file, the Nutrient Data file, Weight file, Footnote file, Nutrient Definition file, and the Data Derivation Code Description file.

Three files contain changes made since SR18 (2005):

- CHG\_FOOD contains records with changes in the descriptive information for a food item.
- CHG\_NUTR contains changes to the following fields: nutrient values, standard errors,

number of data points, source code, and data derivation code.

- CHG\_WGT contains records with changes to the gram weights or measure information.
- CHG\_FTNT contains records with changes to footnotes
- CHG\_NDEF contains records with changes to the nutrient definitions

If the values in any fields have changed, the entire record is included for that file. These files are in the same format as the Food Description, Nutrient Data, Nutrient Definition, Footnote and Weight files. The update files are provided in ASCII format.

Four files contain records that were deleted since SR18 (2005):

- DEL\_FOOD file (Table 15) lists those food items that were deleted from the database.
- DEL\_NUTR file (Table 16) lists those nutrient values that were removed from the database.
- DEL\_WGT contains any gram weights that were removed. These records are in the same format as the Weight file (Table 10).
- DEL\_FTNT contains any footnotes that were removed from the database (Table 17). Starting with SR19, if a given footnote applied to more than one nutrient number, the same footnote number can be used. When these footnote numbers are updated, the extra footnotes are deleted.

**Table 15.—Foods Deleted Format**

Field name	Type	Blank	Description
NDB_No	A 5*	N	Unique 5-digit number identifying deleted item
Shrt_Desc	A 60	N	60-character abbreviated description of food item

\* Index field for Foods Deleted file.

**Table 16. —Nutrients Deleted Format**

Field name	Type	Blank	Description
NDB_No	A 5*	N	Unique 5-digit number identifying the item that contains the deleted nutrient record
Nutr_No	A 3	N	Nutrient number of deleted record

\*Index field for Nutrients Deleted file.

**Table 17.—Footnotes Deleted Format**

Field name	Type	Blank	Description
NDB_No	A 5*	N	Unique 5-digit number identifying the item that contains the deleted nutrient record
Nutr_No	A 3	N	Nutrient number of deleted record
Footnt_Typ	A 1	N	Type of footnote of deleted record

\* Index field for Footnotes Deleted file.

Update files in ASCII are also provided for the Abbreviated file:

- CHG\_ABBR file contains records for food items where a food description, household weight, refuse value, or nutrient value was added, changed, or deleted since SR17. This file is in the same format as the Abbreviated file (Table 14).
- DEL\_ABBR contains food items that were removed from the database; it is in the same format as DEL\_FOOD.
- ADD\_ABBR contains food items added since SR18; it is also in the same format as the Abbreviated file.

## **Summary**

Values for choline and betaine from the USDA Database for the Choline Content of Common Foods (USDA, 2004) have been incorporated into the database, that is SR19. Values for fluoride from the USDA National Fluoride Database of Selected Beverages and Foods - Release 2 (USDA, 2005) have also been incorporated into this update of the database. Other food items, including a number of items for the American Indian/Alaskan Native database, and nutrients/food component values have been updated and expanded as described in “Specific Changes for SR19” (p. 1). SR20, which will be released during mid-2007 will contain additional items and updates.

## References

- Adams, C.F. 1975. Nutritive value of American foods in common units. U.S. Department of Agriculture, Agricultural Handbook 456.
- Association of Official Analytical Chemists. 2003. Official methods of analysis, 17th ed., 2<sup>nd</sup> revision. AOAC International, Gaithersburg, MD.
- Booth, S.L., K.W. Davidson, and J.A. Sadowski. 1994. Evaluation of an HPLC method for the determination of phylloquinone (vitamin K<sub>1</sub>) in various food matrices. *Journal of Agricultural and Food Chemistry* 42:295–300.
- Chun, J., Martin, J.A., Chen, L., Lee, J., Ye, L., Eitenmiller, R. 2004. A differential assay of folic acid and total folate in foods containing enriched cereal-grain products to calculate µg dietary folate equivalents (µg DFE). *Journal of Food Composition and Analysis* 19(2-3):182-187.
- Fulton, L., E. Matthews, and C. Davis. 1977. Average weight of a measured cup of various foods. U.S. Department of Agriculture, Home Economics Research Report 41.
- Gebhardt, S.E., and R.G. Thomas. 2002. Nutritive value of foods. U.S. Department of Agriculture, Home and Garden Bulletin 72.\*
- Holden, J.M., S.A. Bhagwat, and K.Y. Patterson. 2002. Development of a multi-nutrient data quality evaluation system. *Journal of Food Composition and Analysis* 15(4):339–348.\*
- Holden, J.M., A.L. Eldridge, G.R. Beecher, et al. 1999. Carotenoid content of U.S. foods: An update of the database. *Journal of Food Composition and Analysis* 12:169–196.\*
- INFOODS. 2006. Tagnames for Food Components. INFOODS website. [http://www.fao.org/infoods/tagnames\\_en.stm](http://www.fao.org/infoods/tagnames_en.stm).
- Institute of Medicine. 2001. Dietary reference intakes for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium, and zinc. National Academy Press, Washington, DC.
- Institute of Medicine. 2000. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. National Academy Press, Washington, DC.
- Institute of Medicine. 1998. Dietary reference intakes for thiamin, riboflavin, niacin, vitamin B<sub>6</sub>, folate, vitamin B<sub>12</sub>, pantothenic acid, biotin, and choline. National Academy Press, Washington, DC.
- Jones, D.B. 1941. Factors for converting percentages of nitrogen in foods and feeds into percentages of protein. U.S. Department of Agriculture, Circular 83, slight revision.\*

Koc, H., Mar, M. H., Ranasinghe, A., Swenberg, J. A., and Zeisel, S. H. 2002. Quantitation of choline and its metabolites in tissues and foods by liquid chromatography/electrospray ionization-isotope dilution mass spectrometry. *Analytical Chemistry*. 74:4734-4740.

Korz, S., and N.L. Johnson. 1988. Encyclopedia of statistical sciences. Vol. 8, pp. 261–262, John Wiley and Son, New York, NY.

Kubota, J., and W.H. Allaway. 1972. Geographic distribution of trace element problems. *In* J.J. Mortvedt, ed., *Micronutrients in Agriculture: Proceedings of Symposium held at Muscle Shoals, Alabama*, pp. 525–554. Soil Science Society of America, Madison, WI.

Lee, J., Suknark, K., Kluitse, Y., Philips, R.D, Eitenmiller, R.R. 1999. Rapid liquid chromatographic assay of vitamin E and retinyl palmitate in extruded weaning foods. *Journal of Food Science* 64:968-972.

Leonard, S.W., C.K. Good, E.T. Gugger, and M.G. Traber. 2004. Vitamin E bioavailability from fortified breakfast cereal is greater than that from encapsulated supplements. *American Journal of Clinical Nutrition* 79:86–92.

Liem, I.I.H., K.H. Steinkraus, and J.C. Cronk. 1977. Production of vitamin B<sub>12</sub> in tempeh: A fermented soybean food. *Applied Environmental Microbiology* 34:773–776.

Livesay, G., and E. Marinos. 1988. Estimation of energy expenditure, net carbohydrate utilization, and net fat oxidation and synthesis by indirect calorimetry: Evaluation of errors with special reference to the detailed composition of fuels. *American Journal of Clinical Nutrition* 47:608–628.

Mangels, A.R., J.M. Holden, G.R. Beecher, et al. 1993. Carotenoid content of fruits and vegetables: An evaluation of analytic data. *Journal of the American Dietetic Association* 93:284–296.

Martin, J.I., W.O. Landen, A.M. Soliman, and R.R. Eitenmiller. 1990. Application of a tri-enzyme extraction for total folate determination in foods. *Journal of the Association of Analytical Chemistry* 73:805–808.

Matthews, R.H., and Y.J. Garrison. 1975. Food yields summarized by different stages of preparation. U.S. Department of Agriculture, *Agriculture Handbook* 102.\*

Merrill, A.L., and B.K. Watt. 1973. Energy value of foods: Basis and derivation, revised. U.S. Department of Agriculture, *Agriculture Handbook* 74.\*

Murphy, E.W., P.E. Criner, and B.C. Gray. 1975. Comparison of methods for determining retentions of nutrients in cooked foods. *Journal of Agriculture and Food Chemistry* 23:1153.\*

National Academy of Sciences, National Research Council. 1989. Recommended dietary allowances, 10th ed. National Academy Press, Washington, DC.

Pehrsson, P.R., D.B. Haytowitz, J.M. Holden, and C.R. Perry. 2000. USDA's national food and nutrient analysis program: Food sampling. *Journal of Food Composition and Analysis* 13:379–389.\*

Reamer, D.C., and C. Veillon. 1981. Determination of selenium in biological materials by stable isotope dilution gas chromatography-mass spectrometry. *Analytical Chemistry* 53(14):2166–2169.

Schakel, S.F., I.M. Buzzard, and S.E. Gebhardt. 1997. Procedures for estimating nutrient values in food composition databases. *Journal of Food Composition and Analysis* 10:102–114.\*

Sheppard, A.J. 1992. *Lipid Manual: Methodology Suitable for Fatty acid-Cholesterol Analysis*. William C. Brown Publishers, Dubuque, IA.

U.S. Department of Agriculture, Agricultural Research Service. 1976. Composition of foods: Dairy and egg products; raw, processed, prepared. *Agriculture Handbook* 8–1.

U.S. Department of Agriculture, Agricultural Research Service. 1977. Composition of foods: Spices and herbs; raw, processed, prepared. *Agriculture Handbook* 8–2.

U.S. Department of Agriculture, Agricultural Research Service. 1978. Composition of food: Baby foods; raw, processed, prepared. *Agriculture Handbook* 8–3.

U.S. Department of Agriculture, Agricultural Research Service. 1979. Composition of foods: Fats and oils; raw, processed, prepared. *Agriculture Handbook* 8–4.

U.S. Department of Agriculture, Agricultural Research Service. 1979. Composition of foods: Poultry products; raw, processed, prepared. *Agriculture Handbook* 8–5.

U.S. Department of Agriculture, Agricultural Research Service. 1980. Composition of foods: Soups, sauces, and gravies; raw, processed, prepared. *Agriculture Handbook* 8–6.

U.S. Department of Agriculture, Agricultural Research Service. 1980. Composition of foods: Sausages and luncheon meats; raw, processed, prepared. *Agriculture Handbook* 8–7.

U.S. Department of Agriculture, Agricultural Research Service. 1982. Composition of foods: Breakfast cereals; raw, processed, prepared. *Agriculture Handbook* 8–8.

U.S. Department of Agriculture, Agricultural Research Service. 1982. Composition of foods: Fruits and fruit juices; raw, processed, prepared. *Agriculture Handbook* 8–9.

U.S. Department of Agriculture, Agricultural Research Service. 1992. Composition of foods: Pork products; raw, processed, prepared. *Agriculture Handbook* 8–10.

U.S. Department of Agriculture, Agricultural Research Service. 1984. Composition of foods: Vegetables and vegetable products; raw, processed, prepared. Agriculture Handbook 8-11.

U.S. Department of Agriculture, Agricultural Research Service. 1984. Composition of foods: Nut and seed products; raw, processed, prepared. Agriculture Handbook 8-12.

U.S. Department of Agriculture, Agricultural Research Service. 1990. Composition of foods: Beef products; raw, processed, prepared. Agriculture Handbook 8-13.

U.S. Department of Agriculture, Agricultural Research Service. 1986. Composition of foods: Beverages; raw, processed, prepared. Agriculture Handbook 8-14.

U.S. Department of Agriculture, Agricultural Research Service. 1987. Composition of foods: Finfish and shellfish products; raw, processed, prepared. Agriculture Handbook 8-15.

U.S. Department of Agriculture, Agricultural Research Service. 1986. Composition of foods: Legumes and legume products; raw, processed, prepared. Agriculture Handbook 8-16.

U.S. Department of Agriculture, Agricultural Research Service. 1989. Composition of foods: Lamb, veal, and game products; raw, processed, prepared. Agriculture Handbook 8-17.

U.S. Department of Agriculture, Agricultural Research Service. 1992. Composition of foods: Baked products; raw, processed, prepared. Agriculture Handbook 8-18.

U.S. Department of Agriculture, Agricultural Research Service. 1991. Composition of foods: Snacks and sweets; raw, processed, prepared. Agriculture Handbook 8-19.

U.S. Department of Agriculture, Agricultural Research Service. 1989. Composition of foods: Cereal grains and pasta; raw, processed, prepared. Agriculture Handbook 8-20.

U.S. Department of Agriculture, Agricultural Research Service. 1988. Composition of foods: Fast foods; raw, processed, prepared. Agriculture Handbook 8-21.

U.S. Department of Agriculture, Agricultural Research Service. 1990. Composition of foods: Raw, processed, prepared. Agriculture Handbook 8, 1989 Supplement.

U.S. Department of Agriculture, Agricultural Research Service. 1991. Composition of foods: Raw, processed, prepared. Agriculture Handbook 8, 1990 Supplement.

U.S. Department of Agriculture, Agricultural Research Service. 1992. Composition of foods: Raw, processed, prepared. Agriculture Handbook 8, 1991 Supplement.

U.S. Department of Agriculture, Agricultural Research Service. 1993. Composition of foods: Raw, processed, prepared. Agriculture Handbook 8, 1992 Supplement.



U.S. Department of Agriculture, Agricultural Research Service. 1992. Provisional table on the selenium content of foods. USDA, HNIS, PT-109.

U.S. Department of Agriculture. 2003. USDA table of nutrient retention factors, release 5. Nutrient Data Laboratory Home Page at <http://www.ars.usda.gov/nutrientdata>.\*

U.S. Department of Agriculture, Agricultural Research Service. 2004. USDA Database for the Choline Content of Common Foods. Nutrient Data Laboratory website: <http://www.ars.usda.gov/nutrientdata/choline>. \*

U.S. Department of Agriculture, Agricultural Research Service. 2004. USDA Food and Nutrient Database for Dietary Studies (release 2.0) [database]. Food Surveys Research Group website: <http://www.ars.usda.gov/ba/bhnrc/fsrg>.

U.S. Department of Agriculture, Agricultural Research Service. 2005. USDA National Fluoride Database of Selected Beverages and Foods - Release 2. Nutrient Data Laboratory website: <http://www.ars.usda.gov/nutrientdata/fluoride>. \*

U.S. Food and Drug Administration, Department of Health and Human Services. 2004. Food Labeling. Code of Federal Regulations, Title 21, Pt. 101. U.S. Government Printing Office website: <http://www.gpoaccess.gov/cfr/index.html>.

Van Winkle S, Levy S.M., Kiritsy M.C., Heilman J.R., Wefel J.S., Marshall T. 1995. Water and formula fluoride concentrations: Significance for infants fed formula. *Pediatric Dentistry*. 17(4):305-310.

Watt, B.K., and A.L. Merrill. 1963. Composition of foods: Raw, processed, prepared. U.S. Department of Agriculture, Agriculture Handbook 8.

Weihrauch, J.L., and A.S. Chatra. 1994. Provisional table on the vitamin K content of foods, revised. USDA, HNIS, PT-104.\*

Weihrauch, J.L., L.P. Posati, B.A. Anderson, and J. Exler. 1977. Lipid conversion factors for calculating fatty acid content of foods. *Journal of the American Oil Chemists Society* 54:36-40.

Weihrauch, J.L., and J. Tamaki. 1991. Provisional table on the vitamin D content of foods. USDA, HNIS, PT-108.\*

Zeisel, S. H., and Blusztajn, J. K. 1994. Choline and human nutrition. *Annual Reviews of Nutrition*. 14:269-296

Zeisel, S.H., Mar, M.H., Howe, J. C., Holden, J. M. 2003. Concentrations of choline – containing compounds and betaine in common foods. *Journal of Nutrition* 133:1302-1307; Erratum. 2003. *Journal of Nutrition*. 133:2918-2919

---

\* Available on NDL's website: <http://www.ars.usda.gov/nutrientdata>

## Appendix A. Abbreviations Used in Short Descriptions

All purpose	ALLPURP
Aluminum	AL
And	&
Apple	APPL
Apples	APPLS
Applesauce	APPLSAUC
Approximate	APPROX
Approximately	APPROX
Arm and blade	ARM&BLD
Artificial	ART
Ascorbic acid	VIT C
Aspartame	ASPRT
Aspartame-sweetened	ASPRT-SWTND
Baby food	BABYFD
Baked	BKD
Barbequed	BBQ
Based	BSD
Beans	BNS
Beef	BF
Beverage	BEV
Boiled	BLD
Boneless	BNLESS
Bottled	BTLD
Bottom	BTTM
Braised	BRSD
Breakfast	BRKFST
Broiled	BRLD
Buttermilk	BTTRMLK
Calcium	CA
Calorie, calories	CAL
Canned	CND
Carbonated	CARB
Center	CNTR
Cereal	CRL
Cheese	CHS
Chicken	CHICK
Chocolate	CHOC
Choice	CHOIC
Cholesterol	CHOL
Cholesterol-free	CHOL-FREE
Chopped	CHOPD
Cinnamon	CINN
Coated	COATD
Coconut	COCNT

Commercial	COMM
Commercially	COMMLY
Commodity	CMDTY
Composite	COMP
Concentrate	CONC
Concentrated	CONCD
Condensed	COND
Condiment, condiments	CONDMNT
Cooked	CKD
Cottonseed	CTTNSD
Cream	CRM
Creamed	CRMD
Dark	DK
Decorticated	DECORT
Dehydrated	DEHYD
Dessert, desserts	DSSRT
Diluted	DIL
Domestic	DOM
Drained	DRND
Dressing	DRSNG
Drink	DRK
Drumstick	DRUMSTK
English	ENG
Enriched	ENR
Equal	EQ
Evaporated	EVAP
Except	XCPT
Extra	EX
Flank steak	FLANKSTK
Flavored	FLAV
Flour	FLR
Food	FD
Fortified	FORT
French fried	FRENCH FR
French fries	FRENCH FR
Fresh	FRSH
Frosted	FRSTD
Frosting	FRSTNG
Frozen	FRZ
Grades	GRDS
Gram	GM
Green	GRN
Greens	GRNS
Heated	HTD
Heavy	HVY
Hi-meat	HI-MT
High	HI

Hour	HR
Hydrogenated	HYDR
Imitation	IMITN
Immature	IMMAT
Imported	IMP
Include, includes	INCL
Including	INCL
Infant formula	INF FORMULA
Ingredient	ING
Instant	INST
Juice	JUC
Junior	JR
Kernels	KRNLS
Large	LRG
Lean	LN
Lean only	LN
Leavened	LVND
Light	LT
Liquid	LIQ
Low	LO
Low fat	LOFAT
Marshmallow	MARSHMLLW
Mashed	MSHD
Mayonnaise	MAYO
Medium	MED
Mesquite	MESQ
Minutes	MIN
Mixed	MXD
Moisture	MOIST
Natural	NAT
New Zealand	NZ
Noncarbonated	NONCARB
Nonfat dry milk	NFDM
Nonfat dry milk solids	NFDMS
Nonfat milk solids	NFMS
Not Further Specified	NFS
Nutrients	NUTR
Nutrition	NUTR
Ounce	OZ
Pack	PK
Par fried	PAR FR
Parboiled	PARBLD
Partial	PART
Partially	PART
Partially fried	PAR FR
Pasteurized	PAST
Peanut	PNUT

Peanuts	PNUTS
Phosphate	PO4
Phosphorus	P
Pineapple	PNAPPL
Plain	PLN
Porterhouse	PRTRHS
Potassium	K
Powder	PDR
Powdered	PDR
Precooked	PRECKD
Preheated	PREHTD
Prepared	PREP
Processed	PROC
Product code	PROD CD
Propionate	PROP
Protein	PROT
Pudding, puddings	PUDD
Ready-to-bake	RTB
Ready-to-cook	RTC
Ready-to-drink	RTD
Ready-to-eat	RTE
Ready-to-feed	RTF
Ready-to-heat	RTH
Ready-to-serve	RTS
Ready-to-use	RTU
Reconstituted	RECON
Reduced	RED
Reduced-calorie	RED-CAL
Refrigerated	REFR
Regular	REG
Reheated	REHTD
Replacement	REPLCMNT
Restaurant-prepared	REST-PREP
Retail	RTL
Roast	RST
Roasted	RSTD
Round	RND
Sandwich	SNDWCH
Sauce	SAU
Scalloped	SCALLPD
Scrambled	SCRMBLD
Seed	SD
Select	SEL
Separable <sup>1</sup>	
Shank and sirloin	SHK&SIRL
Short	SHRT
Shoulder	SHLDR

Simmered	SIMMRD
Skin	SKN
Small	SML
Sodium	NA
Solids	SOL
Solution	SOLN
Soybean	SOYBN
Special	SPL
Species	SP
Spread	SPRD
Standard	STD
Steamed	STMD
Stewed	STWD
Stick	STK
Sticks	STKS
Strained	STR
Substitute	SUB
Summer	SMMR
Supplement	SUPP
Sweet	SWT
Sweetened	SWTND
Sweetener	SWTNR
Teaspoon	TSP
Thousand	1000
Toasted	TSTD
Toddler	TODD
Trimmed <sup>1</sup>	
Trimmed to <sup>1</sup>	
Uncooked	UNCKD
Uncreamed	UNCRMD
Undiluted	UNDIL
Unenriched	UNENR
Unheated	UNHTD
Unprepared	UNPREP
Unspecified	UNSPEC
Unsweetened	UNSWTND
Variety, varieties	VAR
Vegetable, vegetables	VEG
Vitamin A	VIT A
Vitamin C	VIT C
Water	H2O
Whitener	WHTNR
Whole	WHL
Winter	WNTR
With	W/

Without  
Yellow

WO/  
YEL

---

<sup>1</sup> Removed in short description

## **Appendix B.**

## **Other Abbreviations**

ap	as purchased
ARS	Agricultural Research Service
DFE	dietary folate equivalent
dia	diameter
DRI	Dietary Reference Intakes
fl oz	fluid ounce
FNDDS	USDA Food and Nutrient Database for Dietary Studies
g	gram
INFOODS	International Network of Food Data Systems
IU	international unit
kcal	kilocalorie
kJ	kilojoule
lb	pound
mg	milligram
µg, mcg	microgram
ml	milliliter
NDB	Nutrient Databank
NDBS	Nutrient Databank System
NDL	Nutrient Data Laboratory
NFNAP	National Food and Nutrient Analysis Program
NLEA	Nutrition Labeling and Education Act
oz	ounce
RAE	retinol activity equivalent
RE	Retinol Equivalents
RDA	Recommended Dietary Allowances
SR	USDA National Nutrient Database for Standard Reference