

Table of Fourier Transform Pairs of Energy Signals

| Function name | Time Domain $x(t)$ | Frequency Domain $X(\omega)$ |
|-------------------------|---|--|
| FT | $x(t)$ | $X(\omega) = \int_{-\infty}^{\infty} x(t) e^{-j\omega t} dt = \mathcal{F}\{x(t)\}$ |
| IFT | $x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(\omega) e^{j\omega t} d\omega = \mathcal{F}^{-1}\{X(\omega)\}$ | $X(\omega)$ |
| Rectangle Pulse | $rect\left(\frac{t}{T}\right) = \Pi\left(\frac{t}{T}\right) \equiv \begin{cases} 1 & t \leq T/2 \\ 0 & \text{elsewhen} \end{cases}$ | $T \operatorname{sinc}\left(\frac{T}{2\pi} \omega\right)$ |
| Triangle Pulse | $\Lambda\left(\frac{t}{W}\right) \equiv \begin{cases} 1 - t /W & t \leq W \\ 0 & \text{elsewhen} \end{cases}$ | $W \operatorname{sinc}^2\left(\frac{W}{2\pi} \omega\right)$ |
| Sinc Pulse | $\operatorname{sinc}(Wt) \equiv \frac{\sin(\pi \cdot Wt)}{\pi \cdot Wt}$ | $\frac{1}{W} \operatorname{rect}\left(\frac{\omega}{2\pi W}\right)$ |
| Exponential Pulse | $e^{-a t } \quad a > 0$ | $\frac{2a}{a^2 + \omega^2}$ |
| Gaussian Pulse | $\exp\left(-\frac{t^2}{2\sigma^2}\right)$ | $(\sigma\sqrt{2\pi}) \exp\left(-\frac{\sigma^2 \omega^2}{2}\right)$ |
| Decaying Exponential | $\exp(-at)u(t) \quad \operatorname{Re}\{a\} > 0$ | $\frac{1}{a + j\omega}$ |
| Sinc ² Pulse | $\operatorname{sinc}^2(Bt)$ | $\frac{1}{B} \Lambda\left(\frac{\omega}{2\pi B}\right)$ |







Table of Fourier Transform Pairs of Power Signals

| Function name | Time Domain x(t) | Frequency Domain X(ω) |
|---------------------|--|--|
| FT | $x(t)$ | $X(\omega) = \int_{-\infty}^{\infty} x(t)e^{-j\omega t} dt = \mathcal{F}\{x(t)\}$ |
| IFT | $x(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} X(\omega)e^{j\omega t} d\omega = \mathcal{F}^{-1}\{X(\omega)\}$ | $X(\omega)$ |
| Impulse | $\delta(t)$ | 1 |
| DC | 1 | $2\pi\delta(\omega)$ |
| Cosine | $\cos(\omega_0 t + \theta)$ | $\pi [e^{j\theta} \delta(\omega - \omega_0) + e^{-j\theta} \delta(\omega + \omega_0)]$ |
| Sine | $\sin(\omega_0 t + \theta)$ | $-j\pi [e^{j\theta} \delta(\omega - \omega_0) - e^{-j\theta} \delta(\omega + \omega_0)]$ |
| Complex Exponential | $\exp(j\omega_0 t)$ | $2\pi\delta(\omega - \omega_0)$ |
| Unit step | $u(t) \equiv \begin{cases} 1 & t \geq 0 \\ 0 & t < 0 \end{cases}$ | $\pi\delta(\omega) + \frac{1}{j\omega}$ |
| Signum | $\text{sgn}(t) \equiv \begin{cases} 1 & t \geq 0 \\ -1 & t < 0 \end{cases}$ | $\frac{2}{j\omega}$ |
| Linear Decay | $1/t$ | $-j\pi \text{sgn}(\omega)$ |
| Impulse Train | $\sum_{n=-\infty}^{\infty} \delta(t - nT_s)$ | $\frac{2\pi}{T_s} \sum_{k=-\infty}^{\infty} \delta\left(\omega - k\frac{2\pi}{T_s}\right)$ |
| Fourier Series | $\sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 t}$, where $a_k = \frac{1}{T_0} \int_{T_0} x(t) e^{-jk\omega_0 t} dt$ | $2\pi \sum_{k=-\infty}^{\infty} a_k \delta(\omega - k\omega_0)$ |

Table of Fourier Transforms of Operations

| Operation | FT Property Given $g(t) \Leftrightarrow G(\omega)$ |
|--------------------|---|
| Linearity | $af(t) + bg(t) \Leftrightarrow aF(\omega) + bG(\omega)$ |
| Time Shifting | $g(t - t_0) \Leftrightarrow e^{-j\omega t_0} G(\omega)$ |
| Time Scaling | $g(at) \Leftrightarrow \frac{1}{ a } G\left(\frac{\omega}{a}\right)$ |
| Modulation (1) | $g(t) \cos(\omega_0 t) \Leftrightarrow \frac{1}{2} [G(\omega - \omega_0) + G(\omega + \omega_0)]$ |
| Modulation (2) | $g(t) e^{j\omega_0 t} \Leftrightarrow G(\omega - \omega_0)$ |
| Differentiation | If $f(t) = \frac{dg(t)}{dt}$, then $F(\omega) = j\omega \cdot G(\omega)$ |
| Integration | If $f(t) = \int_{-\infty}^t g(\alpha) d\alpha$, then $F(\omega) = \frac{1}{j\omega} G(\omega) + \pi G(0) \delta(\omega)$ |
| Convolution | $g(t) * f(t) \Leftrightarrow G(\omega) \cdot F(\omega)$, where $g(t) * f(t) \equiv \int_{-\infty}^{\infty} g(\alpha) f(t - \alpha) d\alpha$ |
| Multiplication | $f(t) \cdot g(t) \Leftrightarrow \frac{1}{2\pi} F(\omega) * G(\omega)$ |
| Duality | If $g(t) \Leftrightarrow z(\omega)$, then $z(t) \Leftrightarrow 2\pi g(-\omega)$ |
| Hermitian Symmetry | If $g(t)$ is real valued then $G(-\omega) = G^*(\omega)$ $(G(-\omega) = G(\omega) \text{ and } \angle G(-\omega) = -\angle G(\omega))$ |
| Conjugation | $g^*(t) \Leftrightarrow G^*(-\omega)$ |
| Parseval's Theorem | $P_{avg} = \int_{-\infty}^{\infty} g(t) ^2 dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} G(\omega) ^2 d\omega$ |

Some Notes:

1. There are two similar functions used to describe the functional form $\sin(x)/x$. One is the $\text{sinc}()$ function, and the other is the $\text{Sa}()$ function. We will only use the $\text{sinc}()$ notation in class. Note the role of π in the $\text{sinc}()$ definition:

$$\text{sinc}(x) \equiv \frac{\sin(\pi x)}{\pi x}; \quad \text{Sa}(x) \equiv \frac{\sin(x)}{x}$$

2. The impulse function, aka delta function, is defined by the following three relationships:

- a. Singularity: $\delta(t - t_0) = 0$ for all $t \neq t_0$

- b. Unity area: $\int_{-\infty}^{\infty} \delta(t) dt = 1$

- c. Sifting property: $\int_{t_a}^{t_b} f(t) \delta(t - t_0) dt = f(t_0)$ for $t_a < t_0 < t_b$.

3. Many basic functions do not change under a reversal operation. Other change signs. Use this to help simplify your results.

- a. $\delta(t) = \delta(-t)$ (in general, $\delta(at) \Leftrightarrow \frac{1}{|a|} \delta(t)$)

- b. $\text{rect}(t) = \text{rect}(-t)$

- c. $\Lambda(t) = \Lambda(-t)$

- d. $\text{sinc}(t) = \text{sinc}(-t)$

- e. $\text{sgn}(t) = -\text{sgn}(-t)$

4. The duality property is quite useful but sometimes a bit hard to understand. Suppose a known FT pair $g(t) \Leftrightarrow z(\omega)$ is available in a table. Suppose a new time function $z(t)$ is formed with the same shape as the spectrum $z(\omega)$ (i.e. the function $z(t)$ in the time domain is the same as $z(\omega)$ in the frequency domain). Then the FT of $z(t)$ will be found to be $z(t) \Leftrightarrow 2\pi g(-\omega)$, which says that the F.T. of $z(t)$ is the same shape as $g(t)$, with a multiplier of 2π and with $-\omega$ substituted for t .

An example is helpful. Given the F.T. pair $\text{sgn}(t) \Leftrightarrow 2/j\omega$, what is the Fourier transform of $x(t)=1/t$? First, modify the given pair to $j/2 \text{sgn}(t) \Leftrightarrow 1/\omega$ by multiplying both sides by $j/2$. Then, use the duality function to show that $1/t \Leftrightarrow 2\pi j/2 \text{sgn}(-\omega) = j\pi \text{sgn}(-\omega) = -j\pi \text{sgn}(\omega)$.