



Classifying actions of the alternating group on Riemann surfaces

Jen Paulhus



Abstract group A_{24}

Group information

Description: A_{24}

Order: $310\cdot\cdot\cdot000 = 2^{21} \cdot 3^{10} \cdot 5^4 \cdot 7^3 \cdot 11^2 \cdot 13 \cdot 17 \cdot 19 \cdot 23$

Exponent: $5354228880 = 2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23$

Automorphism group: Group of order $620\cdot\cdot\cdot000 = 2^{22} \cdot 3^{10} \cdot 5^4 \cdot 7^3 \cdot 11^2 \cdot 13 \cdot 17 \cdot 19 \cdot 23$ (generators)

Outer automorphisms: C_2 , of order 2

Composition factors: A_{24}

Derived length: 0

This group is [nonabelian](#) and [simple](#) (hence [nonsolvable](#), [perfect](#), [quasisimple](#), and [almost simple](#)).

Group statistics

Order	1	2	3	4	5
Elements	1	8792390355903	13428028220072048	2542924546378413120	1725747644222610624
Conjugacy classes	1	6	8	18	4
Divisions	1	6	8	18	4
Autjugacy classes	1	6	8	18	4

Minimal Presentations

Permutation degree: 24

Transitive degree: 24

Rank: not computed

Inequivalent generating tuples: not computed



Properties

Label: 310224200866619719680000.a

Order: $2^{21} \cdot 3^{10} \cdot 5^4 \cdot 7^3 \cdot 11^2 \cdot 13 \cdot 17 \cdot 19 \cdot 23$

Exponent: $2^4 \cdot 3^2 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 \cdot 19 \cdot 23$



Simple: yes

$\#G^{\text{ab}}$: 1

$\#Z(G)$: 1

$\#\text{Aut}(G)$: $2^{22} \cdot 3^{10} \cdot 5^4 \cdot 7^3 \cdot 11^2 \cdot 13 \cdot 17 \cdot 19 \cdot 23$

$\#\text{Out}(G)$: 2

Perm deg.: 24

Trans deg.: 24

Rank: not computed

Related objects

- Subgroups
- Extensions
- Supergroups
- As a transitive group

Downloads

- Group to Gap
- Group to Magma

Aaron Wootton

University of Portland


Preprint on arXiv (2025 and updated version yesterday)

Setup

X a compact Riemann surface of genus g with $G = \text{Aut}(X)$ (finite)

The set of orbits of the action is X/G , also a compact Riemann surface.

The natural map $X \rightarrow X/G$ gives us a branched covering branched at r places.



Riemann's Existence Theorem

A finite group G acts on a compact Riemann surface X of genus $g > 1$ if and only if there are elements of the group

$$a_1, b_1, \dots, a_h, b_h, c_1, \dots, c_r$$

which **generate the group**, satisfy the following equation,

$$\prod_{i=1}^h [a_i, b_i] \prod_{j=1}^r c_j = 1_G$$

$$[a, b] = aba^{-1}b^{-1}$$

h is the genus of X/G

Riemann's Existence Theorem

A finite group G acts on a compact Riemann surface X of genus $g > 1$ if and only if there are elements of the group

$$a_1, b_1, \dots, a_h, b_h, c_1, \dots, c_r$$

which generate the group, satisfy the following equation,

$$\prod_{i=1}^h [a_i, b_i] \prod_{j=1}^r c_j = 1_G$$

and so that $m_j = \text{ord}(c_j)$ satisfy the Riemann Hurwitz formula

$$g = 1 + |G|(h - 1) + \frac{|G|}{2} \sum_{j=1}^r \left(1 - \frac{1}{m_j} \right).$$

signature: $(h; m_1, \dots, m_r)$

Riemann's Existence Theorem

A finite group G acts on a compact Riemann surface X of genus $g > 1$ if and only if there are elements of the group

$$a_1, b_1, \dots, a_h, b_h, c_1, \dots, c_r$$

which generate the group, satisfy the following equation,


$$\prod_{i=1}^h [a_i, b_i] \prod_{j=1}^r c_j = 1_G$$

and so that $m_j = \text{ord}(c_j)$ satisfy the Riemann Hurwitz formula

$$g = 1 + |G|(h - 1) + \frac{|G|}{2} \sum_{j=1}^r \left(1 - \frac{1}{m_j} \right).$$

generating vector: $(a_1, b_1, \dots, a_h, b_h, c_1, \dots, c_r)$


Easy to compute.



Potential signatures are those signatures $(h; m_1, \dots, m_r)$ which satisfy the Riemann Hurwitz formula:

$$g = 1 + |G|(h - 1) + \frac{|G|}{2} \sum_{j=1}^r \left(1 - \frac{1}{m_j} \right).$$

Actual signatures are those which also have a generating vector associated to them.



Hard to compute.

Our Question

Which *groups* only have a finite number of potential signatures which fail to be actual signatures?

We say such groups **act with almost all signatures (or are AAS)**.

Carvacho, P., Tucker, Wootton (2021)

Any non-abelian finite simple group is AAS.

Carvacho, P., Tucker, Wootton (2021)

Any non-abelian finite simple group is AAS.

Question

Classify all actions for non-abelian finite simple groups.

$\mathcal{O}(G) = \{\text{Ord}(g) : g \in G\} - \{1\}$ is the order set.

With a very small number of exceptions, any signature of the form

$$(h; \underbrace{n_1, \dots, n_1}_{t_1}, \underbrace{n_2, \dots, n_2}_{t_2}, \dots, \underbrace{n_s, \dots, n_s}_{t_s})$$

for $n_i \in \mathcal{O}(G)$ and $t_i \in \mathbb{Z}^+$ is a **potential signature**.

P., Wootton

For any non-abelian finite simple group G , any potential signature $(h; m_1, \dots, m_r)$ with $h > 1$ and the set of $m_i \in \mathcal{O}(G)$ possibly empty (i.e. no ramification) is an actual signature.

We need to find generators $a_1, b_1, \dots, a_h, b_h, c_1, \dots, c_r$ in G (may be no c_i) so that:

$$\prod_{i=1}^h [a_i, b_i] \prod_{j=1}^r c_j = 1_G.$$

Ore Conjecture

If G is a finite non-abelian simple group, then every element of G is a commutator.

- Ore proved for A_n (1957)
- Thompson proved for $\text{PSL}(2, q)$ (1961-1962)
- Neubüser, Pahlings, Cleavers for the sporadic simple groups (1984)
- Ellers and Gordeev (1998) and Liebeck, O'Brien, Shalev, and Tiep (2010) for simple groups of Lie type

It is well known that finite simple groups may be generated by two elements in the group.

Pick a_h and b_h two generators of the group. If $r \geq 1$, also choose c_i arbitrary elements of the group of order m_i .

Since every element of the group is a commutator, $([a_h, b_h] \cdot c_1 \cdots c_r)^{-1}$ is a commutator, call it $[a_1, b_1]$. If $h > 2$ we also let $a_i = b_i = 1_G$ for $1 < i < h$.

$$\text{Then } \prod_{i=1}^h [a_i, b_i] \prod c_i = 1_G.$$

Question

Classify all actions for non-abelian finite simple groups.

Question

Classify all actions for ~~non-abelian finite~~ ^{the alternating groups} ~~simple groups~~.

Question

Classify all actions for ~~non-abelian finite~~ ^{the alternating groups} ~~simple groups.~~ [^]

with quotient genus > 0 .

Conder (1980)

The signature $(0; 2, 3, 7)$ is the actual signature for the alternating group A_n for all but a finite number of n .

In particular all $n \geq 168$ work. There are 64 cases for $n \leq 167$ where this signature does not act.

P., Wootton

For $n > 6$, every potential signature $(h; m_1, \dots, m_r)$ for $h \geq 1$ is an actual signature for the group A_n .

In A_5 , there is no action with signature $(1; 2)$ and in A_6 there is no action with signature $(1; 3)$.

For $h = 1$ we need to find c_1, \dots, c_r and a, b in A_n so that $aba^{-1}b^{-1}c_1 \cdots c_r = 1_G$, and these elements generate A_n .

Three cases:

- at least 2 branch points (periods)
- 1 branch point $(1; k)$
- $n \leq 40$ (checked via computers)

(can't have no ramification with $h = 1$)

A subgroup G of A_n is **transitive** if every number in $\Omega = \{1, \dots, n\}$ is sent to every other number via some element of the subgroup. In other words, the action of the subgroup on Ω is transitive.

A subgroup G of A_n is **transitive** if every number in $\Omega = \{1, \dots, n\}$ is sent to every other number via some element of the subgroup. In other words, the action of the subgroup on Ω is transitive.

A subset B of Ω is called a **block** if every element of G either fixes B or sends all elements of B outside of B .

A group G is **primitive** if it is transitive and its only blocks are the whole group, the empty set, and sets containing single elements of Ω .

For $h = 1$ we need to find c_1, \dots, c_r and a, b in A_n so that $aba^{-1}b^{-1}c_1 \cdots c_r = 1_G$, and these elements generate A_n .

Three cases:

- at least 2 branch points
- 1 branch point $(1; k)$
- $n \leq 40$ (checked via computers)

- By a result of Bertram (1972), any element of A_n can be written as the product of two cycles of prime length p for $p > 3n/4$ if such a prime exists.

- By a result of Bertram (1972), any element of A_n can be written as the product of two cycles of prime length p for $p > 3n/4$ if such a prime exists.
- There always exists a prime p with $3n/4 < p < n - 2$ for sufficiently large n . So find such prime cycles for the element $(c_1 \cdots c_r)^{-1}$. Call the prime cycles a and b_1 .



We describe how to choose the c_i a few slides from now.

- By a result of Bertram (1972), any element of A_n can be written as the product of two cycles of prime length p for $p > 3n/4$ if such a prime exists.
- There always exists a prime p with $3n/4 < p < n - 3$ for sufficiently large n . So find such prime cycles for the element $(c_1 \cdots c_r)^{-1}$. Call the prime cycles a and b_1 .
- Any two p -cycles (if p is less than $n-1$) are conjugate in A_n so $b_1 = ba^{-1}b^{-1}$ and then $[a, b] \cdot c_1 \cdots c_r = 1_G$.

Miller (1928)

If a transitive group of degree n contains a cycle of prime order p for $n/2 \leq p \leq n - 3$ it must be either the alternating or symmetric group.

For two or more periods, we show that c_1, \dots, c_r can be chosen to give a transitive action. Then by Miller's result, a, b, c_1, \dots, c_r generate all of A_n .

For $h = 1$ we need to find c and a, b in A_n so that $aba^{-1}b^{-1}c = 1_G$, and these elements generate A_n .

Three cases:

- at least 2 branch points
- 1 branch point $(1; k)$
- $n \leq 40$ (checked via computers)

A classical result attributed to Jordan says that a subgroup of S_n which is a **primitive** permutation group and contains a p -cycle for some prime number $p < n - 2$ must be A_n or S_n .

Jones (2013, Corollary 1.3)

Let G be a primitive permutation group of finite degree n , containing a cycle with k fixed points. Then $G \geq A_n$ if $k \geq 3$.

This argument depends on the parity of n . If n is **odd...**

Bertram's result actually says that every element can be written as a product of two ℓ -cycles for any $\lceil \frac{3n}{4} \rceil \leq \ell \leq n$.

We write the element c as a product of two ℓ -cycles for $\ell = n - 4$.

We use these ℓ -cycles (and in a few cases specify the cycle type of c) to prove primitivity.

This argument depends on the parity of n . If n is **even...**

A result of Xu (1969) says that, for n even, every element of A_n can be expressed as a product of two permutations which are both conjugate to $a = (1\ 2)(3\ 4\dots n)$ and whose cycles of length 2 have one point in common.

We write the element c as a product of two elements conjugate to a and then prove primitivity.

For $h = 1$ we need to find c_1, \dots, c_r and a, b in A_n so that $aba^{-1}b^{-1}c_1 \cdots c_r = 1_G$, and these elements generate A_n .

Three cases:

- at least 2 branch points
- 1 branch point $(1; k)$
- $n \leq 40$ (checked via computers)

- Code written in Magma and Python.
- Used *Unity*, a collaborative, multi-institution high-performance computing cluster managed by UMass Amherst Research Computing and Data.
- In many cases, we code some of the constructions in our paper and Bertram's paper to confirm that the given generating vectors work.

Code available on [GitHub](#) (2025)

- Since there are an infinite number of potential signatures for each n , we need to find ways to reduce the number of cases to be checked to a finite number.
 - For k odd, if we know $(1; k)$ is a signature with generating vectors (a, b, c) then $(1; k, k)$ is also a signature with generating vectors (a, b, c^2, c^{-1}) .
 - Similarly k even and (a, b, c, c, c^{-1}) for $(1; k, k, k)$.
- For each n we find examples for $(1; k)$ (and $(1; k, k)$ if k is even) for all $k \in \mathcal{O}(G)$.
- We also show that for $k_1, k_2 \in \mathcal{O}(G)$ we can generate the group with an element of order k_1 and one of order k_2 .

The End